

# **The Parkinson`s disease pain classification system (PDPCS): results from an international mechanism-based classification approach**

Veit Mylius, MD<sup>1,2,3\*</sup>; Santiago Perez Lloret, MD, PhD<sup>4\*</sup>; Rubens Gisbert Cury, MD, PhD<sup>5</sup>;  
Manoel J. Teixeira, MD, PhD<sup>5</sup>; Victor Rossetto Barbosa, MD<sup>5</sup>; Egberto R. Barbosa, MD,  
PhD<sup>5</sup>; Larissa Iulle Moreira, MD<sup>5</sup>; Clarice Listik, MD<sup>5</sup>; Ana Mercia Fernandes M.Sc.<sup>5</sup>; Diogo  
de Lacera Veiga, MD<sup>5</sup>; Julio Barbour, MD<sup>5</sup>; Nathalie Hollenstein, M.Sc.<sup>1</sup>; Matthias Oechsner,  
MD<sup>6</sup>; Julia Walch, MD<sup>2</sup>; Florian Brugger MD<sup>2</sup>, Stefan Hägele-Link, MD<sup>2</sup>; Serafin Beer,  
MD<sup>1</sup>; Alexandra Rizos, MSc<sup>7</sup>; K. Ray Chaudhuri, MD, DSc<sup>7,8,9</sup>; Didier Bouhassira, MD,  
PhD<sup>10</sup>; Jean-Pascal Lefaucheur, MD, PhD<sup>11</sup>; Lars Timmermann, MD<sup>3</sup>; Roman Gonzenbach  
MD, PhD<sup>1</sup>; Georg Kägi, MD<sup>2</sup>; Jens Carsten Möller, MD<sup>3,6</sup>; Daniel Ciampi de Andrade, MD,  
PhD<sup>5</sup>

\* both authors contributed equally

<sup>1</sup>Department of Neurology, Center for Neurorehabilitation, Valens, Switzerland

<sup>2</sup>Department of Neurology, Kantonsspital St. Gallen, Switzerland

<sup>3</sup>Department of Neurology, Philipps University, Marburg, Germany

<sup>4</sup>Biomedical Research Center (CAECIHS-UAI), National Research Council (CONICET),  
Buenos Aires, Argentina

<sup>5</sup> Pain Center, LIM-62, Departamento de Neurologia da Faculdade de Medicina da  
Universidade de Sao Paulo, Hospital das Clínicas, Sao Paulo, Brazil

<sup>6</sup>Parkinson center, Center for Neurological Rehabilitation, Zihlschlacht, Switzerland

<sup>7</sup>King's College Hospital, Parkinson's Foundation Centre of Excellence, London, UK

<sup>8</sup>King's College London, Dept. Basic and Clinical Neuroscience, London, UK

<sup>9</sup>The Maurice Wohl Clinical Neuroscience Institute, London, UK

<sup>10</sup>INSERM U-987, Centre d'Evaluation et de Traitement de la Douleur, Hôpital Ambroise Paré, Assistance Publique - Hôpitaux de Paris, Boulogne-Billancourt and Université Versailles-Saint-Quentin, France.

<sup>11</sup>EA 4391, Unité de Neurophysiologie Clinique, Hôpital Henri Mondor, Assistance Publique - Hôpitaux de Paris, Faculté de Médecine de Créteil, Université Paris-Est-Créteil, France.

### **Text Pages and number of tables and figures**

Text pages (including tables and figures): 38

Tables: 5 Figure: 1 Supplementary tables: 5 Supplementary files: 2

### **Corresponding author:**

Prof. Veit Mylius

Center for Neurorehabilitation

7317 Valens, Switzerland

Phone: 0041813031406

[mylius@med.uni-marburg.de](mailto:mylius@med.uni-marburg.de)

### **Disclosures**

We confirm no conflicts of interest related to the research or the manuscript. The research, abstract, or the manuscript have not been presented previously.

1  
2  
3 **The Parkinson`s disease pain classification system (PDPCS): results**  
4  
5  
6 **from an international mechanism-based classification approach**  
7  
8  
9

10  
11 Veit Mylius, MD<sup>1,2,3\*</sup>; Santiago Perez Lloret, MD, PhD<sup>4\*</sup>; Rubens Gisbert Cury, MD, PhD<sup>5</sup>; Manoel  
12  
13 J. Teixeira, MD, PhD<sup>5</sup>; Victor Rossetto Barbosa, MD<sup>5</sup>; Egberto R. Barbosa, MD, PhD<sup>5</sup>; Larissa Iulle  
14  
15 Moreira, MD<sup>5</sup>; Clarice Listik, MD<sup>5</sup>; Ana Mercia Fernandes M.Sc.<sup>5</sup>; Diogo de Lacerda Veiga, MD<sup>5</sup>;  
16  
17 Julio Barbour, MD<sup>5</sup>; Nathalie Hollenstein, M.Sc.<sup>1</sup>; Matthias Oechsner, MD<sup>6</sup>; Julia Walch, MD<sup>2</sup>;  
18  
19 Florian Brugger MD<sup>2</sup>, Stefan Hägele-Link, MD<sup>2</sup>; Serafin Beer, MD<sup>1</sup>; Alexandra Rizos, MSc<sup>7</sup>; K. Ray  
20  
21 Chaudhuri, MD, DSc<sup>7,8,9</sup>; Didier Bouhassira, MD, PhD<sup>10</sup>; Jean-Pascal Lefaucheur, MD, PhD<sup>11</sup>; Lars  
22  
23 Timmermann, MD<sup>3</sup>; Roman Gonzenbach MD, PhD<sup>1</sup>; Georg Kägi, MD<sup>2</sup>; Jens Carsten Möller, MD<sup>3,6</sup>;  
24  
25 Daniel Ciampi de Andrade, MD, PhD<sup>5</sup>  
26  
27  
28  
29  
30  
31  
32  
33

34 \* both authors contributed equally  
35  
36  
37

38  
39 <sup>1</sup>Department of Neurology, Center for Neurorehabilitation, Valens, Switzerland  
40

41 <sup>2</sup>Department of Neurology, Kantonsspital St. Gallen, Switzerland  
42

43 <sup>3</sup>Department of Neurology, Philipps University, Marburg, Germany  
44

45  
46 <sup>4</sup>**Biomedical Research Center (CAECIHS-UAI), National Research Council (CONICET), Buenos**  
47  
48 **Aires, Argentina**  
49

50  
51 <sup>5</sup> Pain Center, LIM-62, Departamento de Neurologia da Faculdade de Medicina da Universidade de  
52  
53 Sao Paulo, Hospital das Clínicas, Sao Paulo, Brazil  
54

55  
56 <sup>6</sup>Parkinson center, Center for Neurological Rehabilitation, Zihlschlacht, Switzerland  
57

58  
59 <sup>7</sup>King's College Hospital, Parkinson`s Foundation Centre of Excellence, London, UK  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

<sup>8</sup>King's College London, Dept. Basic and Clinical Neuroscience, London, UK

<sup>9</sup>The Maurice Wohl Clinical Neuroscience Institute, London, UK

<sup>10</sup>INSERM U-987, Centre d'Evaluation et de Traitement de la Douleur, Hôpital Ambroise Paré, Assistance Publique - Hôpitaux de Paris, Boulogne-Billancourt and Université Versailles-Saint-Quentin, France.

<sup>11</sup>EA 4391, Unité de Neurophysiologie Clinique, Hôpital Henri Mondor, Assistance Publique - Hôpitaux de Paris, Faculté de Médecine de Créteil, Université Paris-Est-Créteil, France.

### **Text Pages and number of tables and figures**

Text pages (including tables and figures): **38**

Tables: **5** Figure: 1 Supplementary tables: **5** Supplementary files: 2

### **Corresponding author:**

Prof. Veit Mylius

Center for Neurorehabilitation

7317 Valens, Switzerland

Phone: 0041813031406

[mylius@med.uni-marburg.de](mailto:mylius@med.uni-marburg.de)

### **Disclosures**

We confirm no conflicts of interest related to the research or the manuscript. The research, abstract, or the manuscript have not been presented previously.

## Abstract

1  
2  
3 Pain is a common non-motor symptom in patients with Parkinson’s disease (PD) but the correct  
4  
5 diagnosis of the respective cause remains difficult because suitable tools are lacking, so far. We  
6  
7 developed a framework to differentiate PD- from non-PD-related pain and classify PD-related pain  
8  
9 into three groups based on validated mechanistic pain descriptors (nociceptive, neuropathic, or  
10  
11 nociplastic), which encompasses the previously described PD pain types. Severity of PD-related pain  
12  
13 syndromes was scored by ratings of intensity, frequency, and interference with daily living activities.  
14  
15 The PD-Pain Classification System (PD-PCS) was compared with classic pain measures (ie, brief pain  
16  
17 inventory (BPI) and McGill pain questionnaire (MPQ), PDQ-8 quality of life score, MDS-UPDRS  
18  
19 scores, and non-motor symptoms). 159 non-demented PD patients (disease duration 10.2±7.6 years)  
20  
21 and 37 healthy controls were recruited in four centers. PD-related pain was present in 122 patients  
22  
23 (77%), with 24 (15%) suffering one or more syndromes at the same time. PD-related nociceptive,  
24  
25 neuropathic, or nociplastic pain was diagnosed in 87 (55%), 25 (16%), or 35 (22%), respectively. Pain  
26  
27 unrelated to PD was present in 35 (22%) patients. Overall, PD-PCS severity score significantly  
28  
29 correlated with pain’s BPI and MPQ ratings, presence of dyskinesia and motor fluctuations, PDQ-8  
30  
31 scores, depression and anxiety measures. Moderate intra- and inter-rater reliability were observed. The  
32  
33 PD-PCS is a valid and reliable tool for differentiating PD-related pain from PD-unrelated pain. It  
34  
35 detects and scores mechanistic pain subtypes in a pragmatic and treatment-oriented approach, unifying  
36  
37 previous classifications of PD-pain.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

1  
2 Chronic pain (CP) (ie, pain lasting more than three months) affects 18% to 30% of the general  
3  
4 population.[14; 28] In Parkinson's disease (PD), chronic pain is present in 20% of patients at the time  
5  
6 of the diagnosis associated with the early motor stage and affects up to 80% during the course of the  
7  
8 disease.[4; 15; 46] In addition, a Park Pain type has recently been described as one important non-  
9  
10 motor subtype.[40]  
11

12  
13 **Pain in PD has been previously divided** as i. *de novo* pain temporally related to disease onset,  
14  
15 the symptoms of the disease or its treatment (PD-directly related pain); ii. previous chronic pain  
16  
17 aggravated by the disease or its treatment (PD-indirectly related pain); or iii. pain that is neither caused  
18  
19 nor aggravated by the disease (PD-unrelated pain).[35] A myriad of different pain syndromes has been  
20  
21 described in PD, and several classification systems have been proposed.[9] These various pain types  
22  
23 **propositions** were seldom validated[19; 38] or formally tested,[41] which increases the difficulty in  
24  
25 diagnosing and treating pain in PD patients. We aimed at incorporating these previous approaches to  
26  
27 define PD-related pains and to distinguish them from PD-unrelated pains. According to previous  
28  
29 classifications, pain is considered as PD-related when one of the following conditions apply: when  
30  
31 occurring **along** with the first motor symptoms, when occurring/aggravated during the OFF stage,  
32  
33 when occurring simultaneously with choreatic dyskinesia, or when improved by dopaminergic  
34  
35 treatment.[8; 19; 38]  
36  
37  
38  
39  
40  
41  
42

43  
44 In one given disease entity, pain can be caused by different mechanisms, so that pain diagnosis and  
45  
46 treatment is not etiology-driven, but, instead, mechanism-based.[2] In general, there are three main  
47  
48 pain mechanisms of CP that account for most types of pain: nociceptive, neuropathic and nociplastic  
49  
50 pain.[20; 30] In nociceptive pain, nociceptors are activated by mechanical, thermal or mechanical  
51  
52 stimuli related to actual or potential lesion to non-neural tissues. It includes most of the musculoskeletal  
53  
54 (MSK) pain syndromes, such as osteoarthritis, and other inflammatory conditions where tissue lesion  
55  
56 or inflammation predominate. Neuropathic pain is **associated to** a lesion or disease of the peripheral or  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 central somatosensory system, with specific characteristics (tingling, burning or electric-shock-like  
2 sensations) and location (neurologically plausible).[49] Nociplastic pain syndromes comprise  
3 instances where the nociceptive system is overactive without any evidence of somatosensory system  
4 lesion or peripheral activation of nociceptors due to actual or potential tissue damage.[27; 30] Central  
5 sensitization (i.e. increased responsiveness of a sensory neuron to normal or subthreshold inputs),  
6 which is not specific to a single pain type, plays the key role for the sensory gain of the somatosensory  
7 system in nociplastic pain. As depicted above, the proposition of Wasner and Deuschl for PD-related  
8 pains was followed for this validation and nociplastic pain as a third mechanistic descriptor was added  
9 [52], as previously suggested by Marques et al., 2019.[30]  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

22 We have developed a new, **mechanism-based** classification to differentiate PD-related pain  
23 from PD-unrelated pain, with a further characterization of PD-related pain into three subgroups, to  
24 allow pathophysiology-based treatment to be performed.[12; 35]  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2. Methods

### 2.1. Design

This was an international, cross-sectional, multicenter study with a retest validation step.

### 2.2. Patients and consent

Consecutive inpatients and outpatients with or without pain and with the clinical diagnosis of PD according to the criteria of the United Kingdom PD Society Brain Bank were recruited[22] at the Department of Neurology of the Center for Neurorehabilitation in Valens, the Parkinson Center at the Center for Neurological Rehabilitation Zihlschlacht, the Department of Neurology of Kantonsspital St. Gallen in Switzerland and the Hospital das Clínicas of the University of Sao Paulo in Brazil. Age-matched healthy individuals were also included to detect whether the PD-PCS can differentiate PD patients from controls by the pain level. Informed consent was obtained from all subjects before participation in the study. The study protocol was approved by the local Institutional Review Boards in Switzerland (BASEC: 00502) and Brazil (0105/10).

### 2.3. Inclusion and exclusion criteria

Adult PD patients with or without pain that could stay in the ON-state during clinical assessments were included. Participants were screened for potential dementia using the Mini Mental Status Examination (MMSE; exclusion criterion; cut-off < 25). Patients with Deep Brain Stimulation (DBS) and LCIG pump therapy were also excluded.



#### 2.4. Development of the scale

The Parkinson's Disease Pain Classification System (PD-PCS) is a rater-based scale (**Fig. 1 and Suppl. 1**). The generation of the sequence of steps and items were based on formal meeting with pain specialists (doctors, nurses, physiotherapists and psychologists), as well as, movement disorders experts. The scale was based on review articles previously published by the authors.[12; 35] Its main aim was to: i. ascertain that pain is related to PD (irrespective of being directly-or indirectly-related) rather than unrelated to PD; ii. classify the existing pain into one of the three mechanistic descriptors of CP (i.e. nociceptive, neuropathic and nociplastic). A "severity" score was based on the intensity of pain (on a scale from 0 to 10), multiplied by its frequency and the impact in daily living (each using a three-point Likert score) so that scores can range from 0-90 for each pain type.

Within each mechanistic pain descriptor (nociceptive, neuropathic, and nociplastic), classical pain related situations in PD were included, based on classical case descriptions of pain in PD as well as previous tentative proposals to classify PD-related pain.[8; 19; 20; 38; 52] The *douleur neuropathique-4* questionnaire (DN-4) was used to classify pain as neuropathic.[5] We also took into account insights from recent studies suggesting that some particular types of musculoskeletal pain syndromes such as myofascial pain do occur in PD in a prevalent proportion of patients, being particularly responsive to DBS treatment.[11; 12] The scoring system and a detailed discussion on its structure are included (**Suppl. 1 and 2**). **The final three-item-based model has been subject to previous peer review publications [30] and has benefited from presentations in workshops and inputs received at both international movement disorders and pain meetings.[12; 20; 35] Here, only pain-related to PD was analyzed as we assumed that pain unrelated to PD corresponds to pain seen in the general population and since the PD-PCS aimed at assessing PD-related pains.**

## 2.5. Patient assessment

1  
2 PD patients were clinically examined and underwent the UPDRS-III protocol by neurologists  
3 specialized in movement disorders. Raters assessed patient's pain with the classification tool in a  
4  
5 standardized way (Suppl. 2). The PD-PCS was assessed separately for each pain type associated with  
6  
7 PD and the main pain type mentioned in each group (nociceptive, neuropathic, nociplastic) was  
8  
9 documented and analyzed.  
10  
11  
12  
13

14 At baseline, general information concerning PD history was gathered. Medication intake was  
15 recorded (levodopa equivalents were calculated according to Tomlinson et al.[48]). Afterwards,  
16 patients completed the following questionnaires and tests: PD-PCS, Brief Pain Inventory (BPI),[39]  
17 clock-drawing test,[37] QoL in PD questionnaire (PDQ-8),[31] hospital anxiety and depression scale  
18 (HADS),[26] McGill pain questionnaire short-form 1 (MPQ),[32] and the Wearing-off questionnaire-  
19 9 (WOQ-9).[44] Finally, the Movement Disorders Society revision of the unified Parkinson's disease  
20 rating scale parts III and IV (MDS-UPDRS-III and -IV) were evaluated.[24]  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 A part of the questionnaires was reassessed at a second visit after 7 days (5-10 days) by the  
32 same rater to determine intra-rater reliability and simultaneously by a second rater blinded to the  
33 assessments of the first rater to determine inter-rater reliability. Patients were assessed by PD-PCS,  
34 BPI, and MPQ. Additionally, the Clinical Global Impression of Change (CGIC) was assessed by the  
35 patient (Patient's Global Impression of Change (PGIC) and by the physician (Clinical Global  
36 Impression of Change (CGIC)) to determine if patients were stable at the second visit for intra-rater  
37 comparisons.[6]  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

## 2.6. Sample Size

51 By using the 10-times-item rule and counting 8 (PD-PCS) items, 80 PD patients with pain would be  
52 needed. An extra 20% of patients was recruited to account for lost data, thus sample size was adjusted  
53 to 100. As about 60 % of the patients was expected to suffer from pain, 150 patients were included to  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 reach 100 patients with pain. A sub-sample of 40 patients with pain was considered enough for  
2 calculating reliability at a second visit. A sample of 40 non-PD healthy controls, so as to have one  
3 control for about 2 patients with pain, was included.  
4  
5  
6  
7  
8

### 9 *2.7. Data Analyses*

10 Data from the single centers was collected by the leading centers and transferred to the Institute of  
11 Cardiology Research, University of Buenos Aires, National Research Council (CONICET-ININCA),  
12 Buenos Aires, Argentina, for analyses. Comparisons between controls and patients of numerical or  
13 categorical variables were performed by Student's t-test or chi-square test, or their non-parametric  
14 homologues when assumptions were not met.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

### 26 *2.8. Validation analysis*

27  
28  
29 1. Acceptability: proportion of missing data, score distribution, skewness, and floor and ceiling effects  
30 were evaluated. Floor and ceiling effects were **calculated as the proportion** of cases with PD-PCS  
31 scores below 5% or above 95% of total scores, respectively, in patients with pain as assessed by the  
32 BPI;  
33  
34  
35  
36  
37  
38

39 2. Internal consistency was evaluated by Intraclass correlation coefficient (ICC);  
40

41 3. Intra and inter-rater reliability was assessed by Kappa scores for dichotomous variables or ICC for  
42 continuous variables. For these analyzes, only patients identified as "stable" by the CGIC were  
43 included;  
44  
45  
46  
47  
48

49 4. Criterion validity was explored by correlating PD-PCS scores with BPI and MPQ scores using the  
50 Pearson's correlation technique;  
51  
52

53 5. Convergent construct validity was further assessed by correlating the presence and intensity of each  
54 type of pain as assessed by the PD-PCS with MDS-UPDRS part IV, PDQ-8, HADS and WOQ-9;  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

6. Known-group validity was assessed by comparing the scores from the three pain types (subgroups of the PD-PCS) according to QoL and disease characteristics;

7. Internal validity was assessed by a principal component analysis with **non-orthogonal rotation** of pain syndromes intensity scores, as calculated by the PD-PCS.

### 3. Results

#### 3.1. Clinical features and acceptability

One hundred fifty-nine PD patients and 37 healthy controls were recruited in four clinical centers: in Sao Paulo, Brazil (Hospital das Clínicas, Universidade de São Paulo), and in the Eastern part of Switzerland (Center for Neurorehabilitation in Valens, Center for Neurological Rehabilitation Zihlschlacht, and Department of Neurology of Kantonsspital St. Gallen). Main characteristics of patients and control are shown in **Table 1**.

There was no difference in age between groups, but PD patients were more frequently males, less frequently active workers, had higher HADS anxiety and depression scores, and higher clock scores.

Classification of PD-pain was possible for all patients. Assessment with full scale (step 1, step 2 with pain type determination and pain location with the manikin, and step 3 score calculation for the

determined mechanistic pain descriptor) took less than 7 minutes in 85% of cases and less than 10

minutes in the remaining ones. As shown in **Table 1**, 93% of PD patients were affected by pain as

assessed by the BPI vs. 6% of controls ( $p < 0.01$ ). Regarding the PD-PCS, PD-related pain was present

in 122 patients (77%), with 24 (15%) suffering from more than one syndrome at the same time. PD-

related pain with nociceptive, neuropathic, or nociplastic components was diagnosed in 87 (55%), 25

(16%), or 35 (22%) respectively (**the respective pain syndromes are given in Suppl. Table 1**). Most

frequent mixed pain syndromes concerned nociceptive pain combined with nociplastic ( $n = 12, 7\%$ ) or

neuropathic pain ( $n = 9, 6\%$ ). **The pain characteristics according to the DN4 are given for each pain**

**mechanism in Suppl. Table 2.**

Pain unrelated to PD (i.e. neither caused or aggravated by PD) was present in 35 (22%) patients vs. 2 (5%) controls ( $p<0.01$ ).

The number of affected body regions by nociceptive, neuropathic or nociplastic pain was  $4.8 \pm 5.2$ ,  $8.5 \pm 5.8$ , and  $10.1 \pm 8.9$  respectively ( $p<0.01$ ). Affected body regions are shown in **Suppl. Table 3**. In patients with pain according to the BPI, floor effects for nociceptive, neuropathic, nociplastic, and total scores were present in 4%, 4%, 20%, and 32% of cases, respectively. Ceiling effects for these scores in patients with pain were observed in 6%, 0%, 0%, and 0% of cases, respectively. Skewness was 0.98, 2.53, 2.90 and 0.79 for PD-PCS nociceptive, neuropathic, nociplastic, and total scores, respectively. Samples from Brazil and Swiss were similar, apart for the following results: compared to the Swiss, Brazilians were younger ( $60 \pm 12$  vs  $71 \pm 8$   $p<0.05$ ), had higher UPDRS III ( $41.2 \pm 14.9$  vs  $28.9 \pm 12.6$   $p<0.05$ ), and suffered more frequently from LIDs (56% vs 24%,  $p<0.01$ ). Regarding PD-PCS, Swiss patients suffered more frequently from nociplastic pain (33% vs 13%  $p<0.01$ ), and had higher PD-PCS total scores ( $41.2 \pm 14.9$  vs  $28.9 \pm 12.6$   $p<0.05$ ). All other characteristics and results were not different between the samples.

### 3.2. Internal consistency

Consistency of nociceptive, neuropathic, nociplastic severity scores, as assessed by ICC, was 0.08 ( $p=0.90$ ). This confirms that the scale is not unidimensional, as it assessed the presence of different kinds of pain.

### 3.3. Assessment of reliability

Inter- and intra-rater reliability were assessed in patients that came to the second visit and were not considered to exhibit relevant clinical changes in pain as assessed by CGIC. Patients were assessed by the same researcher ( $n=17$ , intra-rater assessment) and by a different one ( $n=24$ , inter-rater assessment). Overall, PD-PCS severity score showed statistically significant intra- (ICC= 0.62) and

inter-rater reliability (ICC=0.59) reliability. Data on reliability of sub-scores are given in **Table 2 and**

**3.**

#### 3.4. Criterion validity

The PD-PCS total score showed significant association with BPI and MPQ scores (**Table 4**).

#### 3.5. Convergent construct validity

Correlations between PD-PCS scores and other variables are shown in **Table 4**. Nociceptive, neuropathic, nociplastic, and total PD-PCS scores correlated with the presence of levodopa-related motor complications, PDQ-8, and HADS anxiety scores. **Nociceptive, neuropathic, and total PD-PCS scores also correlated with HADS depression scores.** Nociceptive, neuropathic, and total scores also correlated with BPI and MPQ scores (**Table 5**).

#### 3.6. Known-group and internal validity

A multinomial logistic regression analysis was used to assess factors associated with pain mechanism (**Suppl. Table 4**). Results showed that nociceptive pain was related to WOQ-9 score (OR, 95% CI= 1.43, 1.15-1.76) and BPI pain score now (1.27, 1.04-1.55); neuropathic pain to WOQ-9 score (1.83, 1.27-2.65), HADS-A score (1.29, 1.01-1.64), BPI pain score now (1.51, 1.13-2.04), and MPQ sensory score (1.19, 1.01-1.40); and nociplastic pain score to WOQ-9 score (1.47, 1.12-1.92). When QoL, through PDQ-8, was stratified according to low (<8), intermediate (9-16), high (17-42) or very high (>42) scores, pain unrelated to PD had a somewhat similar distribution across all strata, while PD-related pain patients were concentrated in the more affected strata. The PD-PCS showed that these differences were even more significantly clear for the nociceptive and neuropathic mechanistic pain descriptors (**Suppl. Table 5**). Correlations between nociceptive, neuropathic, or nociplastic scores with the total PD-PCS scores were all significant (Pearson's  $r=0.74, 0.46, \text{ or } 0.29$ , respectively). A principal

1 component analysis revealed two factors accounting for 73% of variance. Nociceptive and neuropathic  
2 pain loaded positively **each one** in a different factor, while nociplastic pain loaded negatively on **both**  
3 factors.  
4  
5

#### 6 7 8 9 **4. Discussion**

10 We here present an international validation study of a unifying classification system for pain in PD.  
11  
12 This system was able to differentiate PD-related pain from PD-unrelated pain, providing a mechanistic  
13 and treatment-oriented classification of PD-related pain based on pain pathophysiology. This  
14 categorization system showed moderate intra- and inter-rater reliability, **which probably reflects the**  
15 **difficulties of assessing pain in PD.** Moderate correlations with commonly used pain questionnaires  
16 and other scales confirmed criterion and convergent construct validity. On average, patients presented  
17 with moderate pain intensity. One fourth of them had chronic pain that was not directly related to PD,  
18 which is in line with the prevalence of pain in the general population.[14] Pain related to PD was  
19 present in 77% of the sample and comprised a single type of pain in 62%, two types in 14% and three  
20 in 1% of the patients. Interestingly, mixed pain (overlap of neuropathic and nociceptive pain) is  
21 relatively common in the general population,[20] but has never been formally described in PD. The  
22 result that 15% of patients with pain directly related to PD have more than one pain type is clinically  
23 relevant and may impact not only on treatment approaches, but also the design of future trials [20].  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 The PD-PCS score was significantly correlated with those from commonly used questionnaires  
44 such as the BPI and the MPQ. It also showed correlations with QoL, and mood scores. Interestingly,  
45 the total score of the PD-PCS correlated with the MDS-UPDRS-IV score, but not with the motor score.  
46 This finding may be related to the fact that all patients were assessed in the “ON” state. Alternatively,  
47 the dissociation between motor and pain state has been described in several instances[10; 11; 13] and  
48 argue against a unique musclogenetic[43] origin of pain in PD. Our present data suggest that the three  
49 pain types identified by the PD-PCS are actually different pain syndromes, sharing different  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 characteristics and probably reflecting different mechanistic backgrounds and possibly different  
2 responses to treatment. For instance, we found that higher nociceptive pain scores were found in  
3 patients with worse QoL, while this was not true for nociplastic pain. Also, as expected, patients with  
4 nociceptive pain had more localized pain (ie, shoulder) compared to those with nociplastic pain, who  
5 had widespread pain featuring on average twice the number of pain regions in the body than that  
6 reported by nociceptive pain patients. **This finding is in line with the spatially widespread nature of the**  
7 **pain types (central, non-motor off) classified under the nociplastic umbrella compared to regional MSK**  
8 **pain classified as nociceptive.[1; 20]** In general, nociceptive pain was more commonly located in the  
9 trunk and lower back, while neuropathic and nociplastic pain were rather found in the lower limbs and  
10 on both upper and lower limbs, respectively. Finally, visceral pain **may also** be considered a nociplastic  
11 rather than nociceptive pain, in line with a more accepted view in this regard.[3; 50]

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

On the other hand, it is known that PD lowers pain thresholds, so that patients with more severe motor disease **have, in general, more altered** pain sensitivity.[34] Also, it has been repetitively shown that both dopamine replacement therapy,[21; 25; 29; 36; 42] and DBS[9,13] can partially reverse these changes. Pain in PD was originally related to increased muscle rigidity. Indeed, the musculoskeletal origin of pain in PD[43] has been initially put forth, but later evidence challenged this hypothesis, since many patients with severe rigidity do not have pain. Moreover, severity of motor symptoms does not differ between patients with or without pain[23] and pain begins before motor symptoms in a significant proportion of patients.[15; 16] Finally, there is a lack of correlation between motor improvement and pain relief with DBS treatment.[11] These findings speak in favor of a specific role of dopamine as a modulator of sensory and pain processing involved in PD-related pain. However, whether dopamine-related mechanisms are more involved in nociceptive, neuropathic, or nociplastic mechanisms of PD-related pain remain to be determined. In this view, maybe nearly all PD-related pain syndromes may involve dopamine-based dysfunction in the central nervous system (CNS), but they cannot be considered to be what is usually called “central neuropathic pain”. In fact, “central



1 neuropathic pain” is defined by the finding of lesions in the CNS specifically affecting the  
2 somatosensory structures and leading to the occurrence of pain with “neuropathic” characteristics  
3 (tingling, burning, electric shock-like sensations).[49] Dopamine-related mechanisms in PD-related  
4 pain probably go beyond this restrictive view of central neuropathic pain. Therefore, what is usually  
5 called “central PD pain”[41] needs to be urgently revised.[30] In fact, most patients suffering from  
6 “central PD pain” actually have a complex clinical presentation[17; 51], where pain occurs in diverse  
7 areas of the body, in a context of dysphoria, motor restlessness, akathisia and cognitive acceleration,  
8 frequently in association with dopamine oscillation syndrome.[7; 33] Since these patients lack clear  
9 lesion to the somatosensory system, they do not have a “neuropathic” pain syndrome *stricto sensu*, and  
10 more likely present nociplastic pain. **Our data further support this view, as has been previously put  
11 forth by Marques et al., 2019.[30] Indeed, patients with nociplastic pain had more widespread pain  
12 areas and more intense pain burden, usually caused by dopamine agonist withdrawal syndrome,  
13 dopaminergic dysregulation syndrome,[33; 51] non-motor off periods and visceral pain attacks.[45]**

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31 There have been several attempts to classify pain in PD. The first one was Quinn’s pain  
32 classifications, which segregated PD-related from PD-unrelated pain, but had not been organized as a  
33 questionnaire.[38] This first classification proposed that PD-related pain was associated with  
34 fluctuations of the disease and/or dopaminergic treatment. It included pain preceding diagnosis of PD,  
35 off-period pain, painful dystonic spasms, and peak-of-dose pain. We used several of these  
36 characteristics in step 1 of the PD-PCS in order to classify pain as PD-related, and most instances of  
37 pain during motor off-periods were classified as nociceptive in the PD-PCS since in all instances there  
38 are excessive painful contractions conveyed by muscle innervation.[18; 47] The Non-Motor  
39 Symptoms Scale (NMS) proposed an association of pain with PD by an exclusion of further causes  
40 and when pain occurs in the off-stage and improves by dopaminergic treatment.[8] The most common  
41 pain classification by Ford summarized five different forms (musculoskeletal, dystonic, central,  
42 neuropathic, and akathisia[19]) when pain occurs in relation with the cardinal symptoms of PD as well  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 as with akathisia and dystonia. He further suggested to consider the impact of dopaminergic medication  
2 without addressing if pain was PD-related or PD-unrelated. In one recent approach pain was classified  
3 into neuropathic, nociceptive, and miscellaneous pains.[52] Here we used the definition of PD-related  
4 pains based on the classification of Quinn, additionally including the effects of dopaminergic use in  
5 pain.[38], with a further classification of PD-related pains based on the classification of Wasner and  
6 Deuschl.[52]. This allows the distinction between PD related and unrelated pain. To date, there is one  
7 PD pain scale, the King's Parkinson's Disease Pain Scale (KPPS)[9]. It has been validated exclusively  
8 for PD-directly related pain, and proposes seven pain domains (musculoskeletal pain, chronic pain,  
9 fluctuation-related pain, nocturnal pain, oro-facial pain, discoloration/oedema/swelling and radicular  
10 pain). Importantly, in the King's scale only patients with no other etiology for their pain were included.  
11 This is an important issue since up to 30% of the general population have chronic pain, and excluding  
12 all other etiologies of pain would exclude at least a third of PD patients with pain that could potentially  
13 be aggravated by PD (classified here as pain indirectly related to PD). Also, so far, it has not been  
14 shown whether the different KPPS domains constitute actual distinct pain mechanisms or simply sub-  
15 items of larger pain groups.[30] We propose that our approach is comparable to previous classifications  
16 and scales since it provides an umbrella mechanistic classification of pain in PD that can be further  
17 refined into different PD pain types as proposed by Marques 2019 [30] and the King's approach [9].  
18 With a validated classification system, the treatment of pain in PD could in future be based on the  
19 exact subtype of pain, which so far has not been possible, since the existing classifications have either  
20 not been validated or are not mechanism-based.

## 51 5. Conclusions

52 In summary, we presented the validation of a hierarchical approach for the diagnostic classification of  
53 pain in PD in an attempt to unify previous efforts to classify PD-related pain. Based on four questions  
54 the questionnaire establishes a relation of pain with PD before subdividing it into three pain types  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 according to mechanistic descriptors (nociceptive, neuropathic, nociplastic) and providing scores. The  
2 refinement of the characterization of pain in PD should help improve pain in PD patients in a more  
3 pragmatic and symptom-oriented manner.  
4  
5  
6  
7  
8

## 9 **Acknowledgments**

10 The authors thank Parkinson Schweiz for funding this research. The study was also supported by  
11  
12 Zambon, and Mundipharma. The authors have no conflict of interest to declare  
13  
14  
15  
16  
17  
18  
19

## 20 **References**

- 21  
22 [1] Abuhasira R, Zlotnik Y, Horev A, Ifergane G. Fibromyalgia-Like Syndrome Associated with  
23 Parkinson's Disease-A Cohort Study. *J Clin Med* 2019;8(8).  
24  
25  
26  
27 [2] Attal N, Fermanian C, Fermanian J, Lanteri-Minet M, Alchaar H, Bouhassira D. Neuropathic pain:  
28 are there distinct subtypes depending on the aetiology or anatomical lesion? *Pain*  
29 2008;138:343-353.  
30  
31  
32  
33 [3] Aziz Q, Giamberardino MA, Barke A, Korwisi B, Baranowski AP, Wesselmann U, Rief W, Treede  
34 RD, Pain ITftCoC. The IASP classification of chronic pain for ICD-11: chronic secondary  
35 visceral pain. *Pain* 2019;160:69-76.  
36  
37  
38 [4] Beiske AG, Loge JH, Ronningen A, Svensson E. Pain in Parkinson's disease: Prevalence and  
39 characteristics. *Pain* 2009;141:173-177.  
40  
41  
42 [5] Bouhassira D, Attal N, Alchaar H, Boureau F, Brochet B, Bruxelle J, Cunin G, Fermanian J, Ginies  
43 P, Grun-Overdyking A, Jafari-Schluep H, Lanteri-Minet M, Laurent B, Mick G, Serrie A,  
44 Valade D, Vicaud E. Comparison of pain syndromes associated with nervous or somatic lesions  
45 and development of a new neuropathic pain diagnostic questionnaire (DN4). *Pain* 2005;114:29-  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [6] Busner J, Targum SD. The clinical global impressions scale: applying a research tool in clinical practice. *Psychiatry (Edgmont)* 2007;4:28-37.
- [7] Castrioto A, Thobois S, Carnicella S, Maillet A, Krack P. Emotional manifestations of PD: Neurobiological basis. *Movement disorders* 2016;31:1103-1113.
- [8] Chaudhuri KR, Martinez-Martin P, Brown RG, Sethi K, Stocchi F, Odin P, Ondo W, Abe K, Macphee G, Macmahon D, Barone P, Rabey M, Forbes A, Breen K, Tluk S, Naidu Y, Olanow W, Williams AJ, Thomas S, Rye D, Tsuboi Y, Hand A, Schapira AH. The metric properties of a novel non-motor symptoms scale for Parkinson's disease: Results from an international pilot study. *Movement disorders* 2007;22:1901-1911.
- [9] Chaudhuri KR, Rizos A, Trenkwalder C, Rascol O, Pal S, Martino D, Carroll C, Paviour D, Falup-Pecurariu C, Kessel B, Silverdale M, Todorova A, Sauerbier A, Odin P, Antonini A, Martinez-Martin P, Europar, the INMPDSG. King's Parkinson's disease pain scale, the first scale for pain in PD: An international validation. *Movement disorders* 2015;30:1623-1631.
- [10] Ciampi de Andrade D, Lefaucheur JP, Galhardoni R, Ferreira KS, Brandao Paiva AR, Bor-Seng-Shu E, Alvarenga L, Myczkowski ML, Marcolin MA, de Siqueira SR, Fonoff E, Barbosa ER, Teixeira MJ. Subthalamic deep brain stimulation modulates small fiber-dependent sensory thresholds in Parkinson's disease. *Pain* 2012;153:1107-1113.
- [11] Cury RG, Galhardoni R, Fonoff ET, Dos Santos Ghilardi MG, Fonoff F, Arnaut D, Myczkowski ML, Marcolin MA, Bor-Seng-Shu E, Barbosa ER, Teixeira MJ, Ciampi de Andrade D. Effects of deep brain stimulation on pain and other nonmotor symptoms in Parkinson disease. *Neurology* 2014;83:1403-1409.
- [12] Cury RG, Galhardoni R, Fonoff ET, Perez Lloret S, Dos Santos Ghilardi MG, Barbosa ER, Teixeira MJ, Ciampi de Andrade D. Sensory abnormalities and pain in Parkinson disease and its modulation by treatment of motor symptoms. *Eur J Pain* 2016;20:151-165.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [13] Cury RG, Galhardoni R, Teixeira MJ, Dos Santos Ghilardi MG, Silva V, Myczkowski ML, Marcolin MA, Barbosa ER, Fonoff ET, Ciampi de Andrade D. Subthalamic deep brain stimulation modulates conscious perception of sensory function in Parkinson's disease. *Pain* 2016;157:2758-2765.
- [14] Dahlhamer J, Lucas J, Zelaya C, Nahin R, Mackey S, DeBar L, Kerns R, Von Korff M, Porter L, Helmick C. Prevalence of Chronic Pain and High-Impact Chronic Pain Among Adults - United States, 2016. *MMWR Morb Mortal Wkly Rep* 2018;67:1001-1006.
- [15] Defazio G, Berardelli A, Fabbrini G, Martino D, Fincati E, Fiaschi A, Moretto G, Abbruzzese G, Marchese R, Bonuccelli U, Del Dotto P, Barone P, De Vivo E, Albanese A, Antonini A, Canesi M, Lopiano L, Zibetti M, Nappi G, Martignoni E, Lamberti P, Tinazzi M. Pain as a nonmotor symptom of Parkinson disease: evidence from a case-control study. *Arch Neurol* 2008;65:1191-1194.
- [16] Erro R, Picillo M, Vitale C, Amboni M, Moccia M, Santangelo G, Pellecchia MT, Barone P. The non-motor side of the honeymoon period of Parkinson's disease and its relationship with quality of life: a 4-year longitudinal study. *Eur J Neurol* 2016;23:1673-1679.
- [17] Evans AH, Lees AJ. Dopamine dysregulation syndrome in Parkinson's disease. *Curr Opin Neurol* 2004;17:393-398.
- [18] Ferreira-Sanchez MR, Moreno-Verdu M, Cano-de-la-Cuerda R, Fernandez-de-Las-Penas C, Gueita-Rodriguez J, Ortega-Santiago R. Widespread Pressure Pain Hyperalgesia Is Not Related to Pain in Patients with Parkinson's Disease. *Pain Med* 2019.
- [19] Ford B. Pain in Parkinson's disease. *Movement disorders* 2010;25 Suppl 1:S98-103.
- [20] Freynhagen R, Parada HA, Calderon-Ospina CA, Chen J, Rakhmawati Emril D, Fernandez-Villacorta FJ, Franco H, Ho KY, Lara-Solares A, Li CC, Mimenza Alvarado A, Nimmaanrat S, Dolma Santos M, Ciampi de Andrade D. Current understanding of the mixed pain concept: a brief narrative review. *Curr Med Res Opin* 2019;35:1011-1018.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [21] Gerdelat-Mas A, Simonetta-Moreau M, Thalamas C, Ory-Magne F, Slaoui T, Rascol O, Brefel-Courbon C. Levodopa raises objective pain threshold in Parkinson's disease: a RIII reflex study. *J Neurol Neurosurg Psychiatry* 2007;78:1140-1142.
- [22] Gibb WR, Lees AJ. The relevance of the Lewy body to the pathogenesis of idiopathic Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1988;51:745-752.
- [23] Goetz CG, Tanner CM, Levy M, Wilson RS, Garron DC. Pain in Parkinson's disease. *Movement disorders* 1986;1:45-49.
- [24] Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, Martinez-Martin P, Poewe W, Sampaio C, Stern MB, Dodel R, Dubois B, Holloway R, Jankovic J, Kulisevsky J, Lang AE, Lees A, Leurgans S, LeWitt PA, Nyenhuis D, Olanow CW, Rascol O, Schrag A, Teresi JA, van Hilten JJ, LaPelle N. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Movement disorders* 2008;23:2129-2170.
- [25] Granovsky Y. Conditioned pain modulation: a predictor for development and treatment of neuropathic pain. *Curr Pain Headache Rep* 2013;17:361.
- [26] Hinz A, Braehler E. Normative values for the hospital anxiety and depression scale (HADS) in the general German population. *J Psychosom Res* 2011;71:74-78.
- [27] Kosek E, Cohen M, Baron R, Gebhart GF, Mico JA, Rice AS, Rief W, Sluka AK. Do we need a third mechanistic descriptor for chronic pain states? *Pain* 2016;157:1382-1386.
- [28] Leao Ferreira KA, Bastos TR, Andrade DC, Silva AM, Appolinario JC, Teixeira MJ, Latorre MD. Prevalence of chronic pain in a metropolitan area of a developing country: a population-based study. *Arquivos de neuro-psiquiatria* 2016;74:990-998.
- [29] Lim SY, Farrell MJ, Gibson SJ, Helme RD, Lang AE, Evans AH. Do dyskinesia and pain share common pathophysiological mechanisms in Parkinson's disease? *Movement disorders* 2008;23:1689-1695.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [30] Marques A, Attal N, Bouhassira D, Moisset X, Cantagrel N, Rascol O, Durif F, Brefel-Courbon C. How to diagnose parkinsonian central pain? *Parkinsonism Relat Disord* 2019;64:50-53.
- [31] Martinez-Martin P, Benito-Leon J, Alonso F, Catalan MJ, Pondal M, Zamarbide I. Health-related quality of life evaluation by proxy in Parkinson's disease: approach using PDQ-8 and EuroQoL-5D. *Movement disorders* 2004;19:312-318.
- [32] Melzack R. The short-form McGill Pain Questionnaire. *Pain* 1987;30:191-197.
- [33] Michely J, Volz LJ, Barbe MT, Hoffstaedter F, Viswanathan S, Timmermann L, Eickhoff SB, Fink GR, Grefkes C. Dopaminergic modulation of motor network dynamics in Parkinson's disease. *Brain* 2015;138:664-678.
- [34] Mylius V, Brebbermann J, Dohmann H, Engau I, Oertel WH, Moller JC. Pain sensitivity and clinical progression in Parkinson's disease. *Movement disorders* 2011;26:2220-2225.
- [35] Mylius V, Ciampi de Andrade D, Cury RG, Teepker M, Ehrt U, Eggert KM, Beer S, Kesselring J, Stamelou M, Oertel WH, Moller JC, Lefaucheur JP. Pain in Parkinson's Disease: Current Concepts and a New Diagnostic Algorithm. *Mov Disord Clin Pract* 2015;2:357-364.
- [36] Nandhagopal R, Troiano AR, Mak E, Schulzer M, Bushnell MC, Stoessl AJ. Response to heat pain stimulation in idiopathic Parkinson's disease. *Pain Med* 2010;11:834-840.
- [37] Nishiwaki Y, Breeze E, Smeeth L, Bulpitt CJ, Peters R, Fletcher AE. Validity of the Clock-Drawing Test as a screening tool for cognitive impairment in the elderly. *Am J Epidemiol* 2004;160:797-807.
- [38] Quinn NP, Koller WC, Lang AE, Marsden CD. Painful Parkinson's disease. *Lancet* 1986;1:1366-1369.
- [39] Radbruch L, Loick G, Kiencke P, Lindena G, Sabatowski R, Grond S, Lehmann KA, Cleeland CS. Validation of the German version of the Brief Pain Inventory. *J Pain Symptom Manage* 1999;18:180-187.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [40] Sauerbier A, Jenner P, Todorova A, Chaudhuri KR. Non motor subtypes and Parkinson's disease. *Parkinsonism Relat Disord* 2016;22 Suppl 1:S41-46.
- [41] Schestatsky P, Kumru H, Valls-Sole J, Valldeoriola F, Marti MJ, Tolosa E, Chaves ML. Neurophysiologic study of central pain in patients with Parkinson disease. *Neurology* 2007;69:2162-2169.
- [42] Slaoui T, Mas-Gerdelat A, Ory-Magne F, Rascol O, Brefel-Courbon C. [Levodopa modifies pain thresholds in Parkinson's disease patients]. *Rev Neurol (Paris)* 2007;163:66-71.
- [43] Snider SR, Fahn S, Isgreen WP, Cote LJ. Primary sensory symptoms in parkinsonism. *Neurology* 1976;26:423-429.
- [44] Stacy MA, Murphy JM, Greeley DR, Stewart RM, Murck H, Meng X. The sensitivity and specificity of the 9-item Wearing-off Questionnaire. *Parkinsonism Relat Disord* 2008;14:205-212.
- [45] Storch A, Schneider CB, Wolz M, Sturwald Y, Nebe A, Odin P, Mahler A, Fuchs G, Jost WH, Chaudhuri KR, Koch R, Reichmann H, Ebersbach G. Nonmotor fluctuations in Parkinson disease: severity and correlation with motor complications. *Neurology* 2013;80:800-809.
- [46] Tinazzi M, Del Vesco C, Fincati E, Ottaviani S, Smania N, Moretto G, Fiaschi A, Martino D, Defazio G. Pain and motor complications in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2006;77:822-825.
- [47] Tinazzi M, Recchia S, Simonetto S, Tamburin S, Defazio G, Fiaschi A, Moretto G, Valeriani M. Muscular pain in Parkinson's disease and nociceptive processing assessed with CO2 laser-evoked potentials. *Movement disorders* 2010;25:213-220.
- [48] Tomlinson CL, Stowe R, Patel S, Rick C, Gray R, Clarke CE. Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Movement disorders : official journal of the Movement Disorder Society* 2010;25:2649-2653.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [49] Treede RD, Jensen TS, Campbell JN, Cruccu G, Dostrovsky JO, Griffin JW, Hansson P, Hughes R, Nurmikko T, Serra J. Neuropathic pain: redefinition and a grading system for clinical and research purposes. *Neurology* 2008;70:1630-1635.
- [50] Treede RD, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, Cohen M, Evers S, Finnerup NB, First MB, Giamberardino MA, Kaasa S, Korwisi B, Kosek E, Lavand'homme P, Nicholas M, Perrot S, Scholz J, Schug S, Smith BH, Svensson P, Vlaeyen JWS, Wang SJ. Chronic pain as a symptom or a disease: the IASP Classification of Chronic Pain for the International Classification of Diseases (ICD-11). *Pain* 2019;160:19-27.
- [51] Warren N, O'Gorman C, Lehn A, Siskind D. Dopamine dysregulation syndrome in Parkinson's disease: a systematic review of published cases. *J Neurol Neurosurg Psychiatry* 2017;88:1060-1064.
- [52] Wasner G, Deuschl G. Pains in Parkinson disease--many syndromes under one umbrella. *Nat Rev Neurol* 2012;8:284-294.

## Legends

Fig. 1. The PD-Pain Classification System (PD-PCS) with a complementary QR code for a web-based on-line version.

Table 1. Pain in healthy controls and in PD patients

Table 2: PD-PCS scores and other pain measures in PD patients assessed on two occasions 7 days apart (n=48)

Table 3: Intra- and inter-rater reliability

Table 4. Correlations between PD-PCS scores and other variables at Visit 1 in PD patients (n=159).

Table 5. Correlation of changes in PD-PCS scores with other pain measures

## SDC

Suppl. 1. The PD-Pain Classification System (PD-PCS) questionnaire

Suppl. 2. Instructions for the Parkinson's Disease Pain Classification System (PD-PCS)

Suppl. Table 1. Prevalence of nociceptive, neuropathic and nociplastic pains with respect to the defined pain syndrome at the first visit (n=159)

Suppl. Table 2. Frequency of responses in the DN4 concerning the pain mechanisms at the first visit (n=159)

Suppl. Table 3. Regions affected by types of PD-related pain

Suppl. Table 4. Differences in patients' characteristics according to the type of PD-related pain

Suppl. Table 5. PD-PCS assessment according to PDQ-8

**Funding** The study was supported by Parkinson Schweiz, Zambon, and Mundipharma.

**Competing interests** VM received honoraria from LicherMT, Boston Scientific and Abbvie. SPL received honoraria from IPMDS and consulted for Merz Pharmaceuticals. RGC, MJT, VRB, ERB, LIM, CL, AMF, DLV, JB, NH, MO, JW, SHL, SB, DB, JPL, and JCM report no disclosures. FB received research grants from Baasch-Medicus Foundation and Forschungskommission Kantonsspital St. Gallen, and a travel grant from Abbvie Switzerland. AR received salary support from the Institute for National Health Research (NIHR) Clinical Research Network (CRN) South London and honoraria from Britannia Pharmaceuticals Ltd. KRC received honoraria for advisory boards by AbbVie, UCB, Pfizer, Jazz Pharma, GKC, Bial, Cynapsus, Novartis, Lobsor, Stada, Medtronic, Zambon, Profile, Sunovion, Roche, Therevance, Scion; Honoraria for lectures by AbbVie, Britannia, UCB, Mundipharma, Zambon, Novartis, Boeringer Ingelheim, Neuroderm, Sunovion; Grants (Investigator Initiated) by Britania Pharmaceuticals, AbbVie, UCB, GKC, Bial; Academic grants by EU (Horizon 2020), IMI, Parkinson's UK, NIHR, PDNMG, Kirby Laing Foundation, NPF, MRC. LT received payments as a consultant for Boston Scientific, honoraria as a speaker on symposia sponsored by UCB, Desitin, Boston Scientific, Abott. RG received honoraria from Almirall, Bayer, Sandoz, Biogen, Roche, Novartis, Sanofi. GK received honoraria from Bayer. DCA consulted for Merk, Grunenthal, Cristalia, Novartis. Recipient of Investigator Initiated Research grant from Grunenthal, Mundipharma, Pfizer, Abbot.

**An original** tool to distinguish Parkinson's disease related from non-Parkinson's disease related pain and to classify it into three mechanistic pain descriptors.

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

# PD-Pain Classification System



"QR code" for accessing the on-line application for the PS-PCS





Table 1. Pain in healthy controls and in PD patients

	Healthy controls (n=37)	PD (n=159)	p-value
Males	16 (43%)	99 (62%)	0.04
Age (ys)	65.0±11.5	65.1±11.6	0.96
Right handedness	35 (95%)	156 (98%)	0.22
Married	24 (65%)	112 (70%)	0.51
Active	13 (35%)	13 (8%)	<0.01
PD duration (ys)	-	10.2±7.6	-
MDS-UPDRS III score	-	35.5±15.2	-
MDS-UPDRS IV score	-	6.1±4.6	-
LIDs	-	66 (42%)	-
Daily % score	-	0.6±0.9	-
OFF-time	-	113 (72%)	-
Daily % score	-	1.1±1.0	-
WOQ-9 score	-	4.8±2.6	-
Clock score	1.0±0.1	2.8±1.5	<0.01
PDQ-8 score	-	28.2±23.5	-
HADS-A score	2.7±2.9	7.5±4.1	<0.01
HADS-D score	1.4±2.2	7.4±4.5	<0.01
<b>Antiparkinsonian drugs</b>			
Levodopa	-	146 (92%)	
Agonists	-	75 (47%)	
Other	-	86 (54%)	
Levodopa equivalent dose	-	1050.69±634.78	

**Pain**

Pain reported at BPI	2 (6%)	148 (93%)	<0.01
Maximum pain score	-	7.2±2.6	-
Minimum pain score	-	1.7±2.2	-
Average pain score	-	5.1±2.3	-
Ongoing pain score	-	2.9±2.9	-
MPQ sensory	-	13.3±7.8	-
MPQ affective	-	5.1±4.4	-
MPQ total	-	18.5±11.4	-

**PD-PCS**

No pain		37 (23%)	
PD-unrelated pain	2 (5%)	35 (22%)	<0.01
PD-related pain		122 (77%)	
PD-related pain component:			
Nociceptive	0	87 (55%)	<0.01
Score	0	22.6±29.1	<0.01
Neuropathic	0	25 (16%)	<0.01
Score	0	7.3±19.1	<0.01
Nociplastic	0	35 (22%)	<0.01
Score	0	6.0±16.4	<0.01
One component	-	98 (62%)	-
Two components	-	22 (14%)	-
Three components	-	2 (1%)	-
PD-PCS total score	0	36.0±35.1	<0.01

---

Means±standard deviations are shown. Comparisons were performed by means of chi-squared or t-tests.

PD: Parkinson's Disease; MDS-UPDRS III-IV: Movement Disorders Society Revision of the Unified Parkinson's Disease Rating Scale parts III and IV; LIDs: levodopa-induced dyskinesia; WOQ-9: Wearing-off questionnaire-9; PDQ-8: Quality of life in Parkinson's Disease questionnaire; HADS-A and HADS-D: Hospital Anxiety and Depression scale - Anxiety and depression scores; BPI: Brief Pain Inventory; MPQ: McGill Pain Questionnaire; PD-PCS: Parkinson's Disease - Pain Classification System



Table 2: PD-PCS scores and other pain measures in PD patients assessed on two occasions 7 days apart (n=48)

	Visit 1	Visit 2	$\Delta$ V2-V1	p-value
<b>PD-PDS</b>				
PD-unrelated	8 (17%)	5 (11%)	-	0.45
Nociceptive pain	27 (56%)	34 (71%)	-	0.09
Score	26.5 $\pm$ 30.0	30.2 $\pm$ 28.3	4.2 $\pm$ 26.1	0.30
Neuropathic pain	11 (23%)	5 (10%)	-	0.07
Score	9.5 $\pm$ 20.2	4.6 $\pm$ 14.2	-5.1 $\pm$ 20.7	0.09
Nociplastic pain	9 (19%)	13 (27%)	-	0.22
Score	3.8 $\pm$ 12.7	7.1 $\pm$ 21.7	3.3 $\pm$ 22.1	0.14
PD-PDS total score	40.6 $\pm$ 34.3	43.9 $\pm$ 32.2	3.3 $\pm$ 30.2	0.39
<b>Brief Pain Inventory</b>				
Maximum pain	7.9 $\pm$ 2.0	7.5 $\pm$ 2.2	-0.4 $\pm$ 2.5	0.35
Minimum pain	1.4 $\pm$ 1.6	1.8 $\pm$ 2.0	0.4 $\pm$ 2.1	0.18
Average pain	5.5 $\pm$ 2.0	5.7 $\pm$ 2.4	0.2 $\pm$ 2.3	0.65
Ongoing pain	2.7 $\pm$ 2.8	3.0 $\pm$ 2.9	0.2 $\pm$ 3.3	0.68
<b>McGill Pain Questionnaire</b>				
Sensory	14.6 $\pm$ 7.6	14.6 $\pm$ 8.2	-0.1 $\pm$ 6.5	0.91
Affective	6.1 $\pm$ 4.4	6.6 $\pm$ 4.7	0.6 $\pm$ 3.1	0.21
Total score	20.7 $\pm$ 11.0	21.1 $\pm$ 12.1	0.4 $\pm$ 8.1	0.72
<b>Change scores</b>				
PGIC	-	3.3 $\pm$ 1.4	-	-
Improvement	-	26 (54%)	-	-
No change	-	17 (35%)	-	-

	Visit 1	Visit 2	$\Delta$ V2-V1	p-value
Worsening	-	5 (10%)	-	-
CGIC	-	3.4 $\pm$ 1.1	-	-
Improvement	-	23 (48%)	-	-
No change	-	20 (42%)	-	-
Worsening	-	5 (10%)	-	-

Means $\pm$ standard deviations are shown. Numerical variables were compared by paired t-test and the categorical ones by McNemar test.

PD: Parkinson's Disease; PD-PCS: Parkinson's Disease - Pain Classification System; PGIC: Patient's Global Impression of Change; CGIC: Clinical Global Impression of Change.

Table 3: Intra- and inter-rater reliability

	Intra-rater (n=17)	Inter-rater (n=24)
Nociceptive pain <sup>a</sup>	0.60*	0.40*
Nociceptive Score <sup>b</sup>	0.37*	0.65*
Neuropathic pain <sup>a</sup>	0.43*	0.33
Neuropathic Score <sup>b</sup>	0.34*	0.69*
Nociplastic pain <sup>a</sup>	0.43*	0.23
Nociplastic Score <sup>b</sup>	0.50*	0.04
PD-PCS total score <sup>b</sup>	0.62*	0.59*

Kappa scores (a) or Intraclass Correlation Coefficients (b) are shown (\* p<0.05). Only patients with no change on Clinical Global Impression of Change (CGIC) and Patient's Global Impression of Change (PGIC) were selected for these analyses.

PD-PCS: Parkinson's Disease - Pain Classification System.

Table 4: Correlations between PD-PCS scores and other variables at Visit 1 in PD patients (n=159).

	Nociceptive score	Neuropathic score	Nociplastic score	PD-PCS Total score
MDS-UPDRS-III score	0.08	0.13	-0.07	0.10
MDS-UPDRS-IV score	0.22**	0.04	0.15	0.28**
LIDs daily %	0.18**	0.07	-0.02	0.18**
OFF-time daily %	0.12	0.07	0.10	0.19**
WOQ-9 score	0.06	0.18**	0.20**	0.27**
Clock score	0.03	0.02	0.00	0.04
PDQ-8 score	0.24**	0.18**	0.16**	0.39**
HADS-A	0.25**	0.19**	0.16**	0.40**
HADS-D	0.22**	0.18**	0.05	0.33**
BPI worst	0.33**	0.16**	0.03	0.40**
BPI weakest	0.25**	0.06	-0.04	0.22**
BPI average	0.31**	0.16**	0.16**	0.43**
BPI now	0.28**	0.18**	-0.01	0.32**
MPQ sensory	0.31**	0.35**	0.08	0.49**
MPQ affective	0.31**	0.22**	0.07	0.43**
MPQ total	0.33**	0.32**	0.08	0.50**

Pearson correlation coefficients are shown (\* p<0.05, \*\* p<0.01).

PD-PCS: Parkinson's Disease - Pain Classification System; MDS-UPDRS III-IV: Movement Disorders Society Revision of the Unified Parkinson's Disease Rating Scale parts III and IV; LIDs: levodopa-induced dyskinesia; WOQ-9: Wearing-off questionnaire-9; PDQ-8: Quality of life in Parkinson's Disease questionnaire; HADS-A and HADS-D: Hospital Anxiety and Depression scale - Anxiety and depression scores; BPI: Brief Pain Inventory; MPQ: McGill Pain Questionnaire.

Table 5. Correlation of changes in PD-PCS scores with other pain measures

	Nociceptive score	Neuropathic score	Nociplastic score	Total
BPI worst	0.13	-0.06	0.14	0.26
BPI weakest	-0.01	0.10	-0.16	-0.05
BPI average	0.21	-0.09	-0.20	0.01
BPI now	0.37**	-0.22	-0.27	-0.01
MPQ sensory	0.24	0.10	0.03	0.30**
MPQ affective	0.17	-0.22	0.19	0.17
MPQ total	0.26	0.00	0.09	0.31**

Pearson correlation coefficient are shown. \*  $p < 0.05$ , \*\*  $p < 0.01$ .

PD-PCS: Parkinson's Disease - Pain Classification System; BPI: Brief Pain Inventory;

MPQ: McGill Pain Questionnaire.