

HISTAMINE H4 RECEPTOR EXPRESSION IN TRIPLE NEGATIVE BREAST CANCER: AN EXPLORATORY STUDY.

*Daniela Speisky¹, Mónica Taquez Delgado², Alejandro Iotti¹, Melisa B. Nicoud²,
Ignacio Ospita², Félix Vigovich¹, Pablo Dezanzo¹, Glenda Ernst³, Juan Luis
Uriburu⁴, Vanina A. Medina^{2*}*

(1) Pathology Department, British Hospital, Buenos Aires, Argentina.

*(2) Laboratory of Tumor Biology and Inflammation. Institute for Biomedical
Research (BIOMED), School of Medical Sciences, Pontifical Catholic University
of Argentina (UCA), and the National Scientific and Technical Research Council
(CONICET), Buenos Aires, Argentina.*

(3) Scientific Committee, British Hospital, Buenos Aires, Argentina

(4) Mastology Department, British Hospital, Buenos Aires, Argentina

**Correspondence to:*

*Vanina A. Medina, Ph.D, Laboratory of Tumor Biology and Inflammation.
Institute for Biomedical Research (BIOMED), School of Medical Sciences,
Pontifical Catholic University of Argentina (UCA), and the National Scientific and
Technical Research Council (CONICET), Buenos Aires, Argentina. e-mail:
vanina_medina@uca.edu.ar. Phone: +54-11-4349-0200 ext. 6091*

Running title: H4R in triple negative breast cancer

ABSTRACT

Triple-negative breast cancer (TNBC) is an aggressive breast cancer subtype. There are neither universally accepted prognostic markers, nor molecular targets related to TNBC. The histamine H4 receptor (H4R) has been characterized in TNBC experimental models, demonstrating its critical role in tumor development and progression. In this study, H4R expression was compared in breast cancer subtypes and correlated with clinical features using The Cancer Genome Atlas data (TCGA, Pan-Cancer Atlas). The H4R status was further evaluated by immunohistochemistry in 30 TNBC human samples in relation to clinicopathological parameters. Results indicate that H4R was downregulated in basal-like/TNBC compared with luminal A and normal breast-like tumors. The higher expression of H4R was associated with improved progression free and overall survival outcomes in basal-like/TNBC. H4R immunoreactivity was detected in about 70% of tumors, and its expression was positively correlated with the levels in the histologically normal peritumoral tissue. High H4R expression in peritumoral tissue correlated with reduced number of lymph node involvement and unifocal TNBC while it was associated with increased patient survival. In conclusion, the H4R might represent a potential prognostic biomarker in TNBC. Further studies in large cohorts are needed to better understand the significance of H4R in breast cancer biology.

KEYWORDS: histamine H4 receptor, triple negative breast cancer, prognostic marker, metastasis, immunohistochemistry

INTRODUCTION

Breast cancer is the most frequently diagnosed neoplasia and the leading cause of cancer-related mortality among women worldwide.^{1,2} These tumors are heterogeneous, and present distinct histopathological patterns and clinical behavior. Different molecular subtypes of breast cancer with distinctive biological features have been identified, based on gene expression profiles of human tumors. They include luminal A, luminal B, basal-like, normal breast-like, and human epidermal growth factor receptor 2 (HER2) positive subgroups with different incidence and prognosis.³

Within the pathology-based triple negative tumors, the vast majority fall into the basal-like molecularly classified subtype (around 80%, depending on the study). Triple-negative breast cancer (TNBC) accounts for about 10-20% of all breast cancers and it is considered the most aggressive subtype, lacking estrogen receptor (ER), progesterone receptor (PR), and HER2.⁴⁻⁹ It is associated with poor prognostic features including higher nuclear grade, increased incidence of metastases, and a short recurrence-free interval. Furthermore, there are neither universally accepted prognostic markers to predict outcomes, nor well-defined molecular targets in TNBC subtype.^{3,5,6,8} Therefore, there is an urgent need to establish prognostic factors and to improve TNBC treatments, focusing on the development of novel biomarkers to identify potential patients for a personalized therapeutic approach.⁵⁻⁹ In this regard, one of the most important conditions is an adequate characterization of the tumors and the understanding of the mechanisms involved in TNBC heterogeneity.

The histaminergic system is one of the most interesting and complex biological pathways involved in cancer disease. High histamine biosynthesis and content together with histamine receptors have been reported in different tumors, including gastric, colorectal, esophageal, oral, pancreatic, liver, lung, skin, blood, and breast cancers.^{10,11} The histamine H4 receptor (H4R) was discovered two decades ago, and it has contributed to a better understanding of the histamine roles in health and disease, opening new perspectives in neoplastic research.¹⁰⁻¹² In breast cancer and particularly in TNBC, H4R expression has been well characterized in different *in vitro* and *in vivo* experimental models, demonstrating its critical role in the regulation of tumor proliferation, development, and progression. The administration of histamine or H4R agonists diminished the tumor growth in both immune-deficient and immune-competent TNBC preclinical experimental models.¹²⁻¹⁶ The analysis of TCGA (The Cancer Genome Atlas) data showed that the H4R gene expression is impaired in primary tumors compared with normal tissue in different cancer types.^{11,17} However, the immunohistochemical expression of H4R in TNBC and its prognostic value is completely unknown.

In the present exploratory work, we first compared the H4R expression in breast cancer subtypes using publicly available TCGA data, and correlated H4R mRNA expression with clinical attributes. We corroborated transcriptomic data by analyzing the H4R status in TNBC human samples in relation to clinicopathological parameters. This study will improve the knowledge of the role of H4R in breast cancer progression and could provide a venue for the development of a new diagnostic tool and/or therapeutic target, particularly for those subtypes of breast cancer with limited therapeutic options.

MATERIALS AND METHODS

In silico data analysis

The cBioPortal for Cancer Genomics is an open-access resource for interactive exploration of multidimensional cancer genomics data sets.^{18,19}

Mutations and DNA copy-number data, mRNA expression data, and de-identified clinical and survival data were extracted from cBioPortal employing the TCGA breast cancer (BRCA) PanCancer Atlas data set (n= 1072, 12 male patients were excluded for the analyses) (<http://www.cbioportal.org/>).

Correlations between breast cancer patient survival and *H4R* expression (probe set: 221170_at) were further analyzed by KM plotter, mRNA gene chip (<http://kmplot.com>).²⁰ “Auto select best cutoff” and all datasets were chosen in the analysis. Patient cohorts with high and low H4R expressing tumors were compared by a Kaplan–Meier survival plot, and the hazard ratio with 95% confidence intervals and log-rank P-value were calculated.

Patient selection

Thirty women patients with TNBC that underwent breast surgery at British Hospital of Buenos Aires, Argentina between January 2005 and December 2013 were retrospectively studied using archived paraffin-embedded tumor tissue specimens. The clinical, demographic, and histopathologic data recorded is described in Table 1. Survival data were available for twenty-three patients in a period of 24 months and during that period, five patients died due to breast cancer and eighteen of them were alive. The follow up was not available in

seven patients. A great majority of the patients (n=25; 83.3%) underwent additional therapies (adjuvant chemotherapy and/or radiotherapy).

Male breast cancer, benign lesions and non-epithelial breast tumors were excluded. Poorly preserved samples with extensive necrosis were not used in this study. The study was conducted in accordance with the Declaration of Helsinki and approved by the institutional review board of the British Hospital (CRIHB #925).

Cell culture and immunostaining

The human MDA-MB-231 TNBC and MCF-7 luminal breast cancer cells and HEK293 cells (human cell line originally derived from human embryonic kidney cells) (American Type Tissue Culture Collection, VA, USA) were cultured in RPMI 1640 supplemented with 10% v/v FBS, 0.3 g L⁻¹ glutamine and 0.04 g L⁻¹ gentamicin (Gibco BRL, NY, USA). Cells were maintained at 37°C in a humidified atmosphere containing 5% CO₂. The procedures were previously described.^{12,21} Briefly, cells were cultured on glass coverslips into 12-well plates for 24 hours, and they were then fixed with 4% formaldehyde and endogenous peroxidase activity was blocked with 3% hydrogen peroxide (v/v) in distilled water. After blocking, cells were incubated overnight in a humidified chamber at 4°C with primary rabbit anti-H4R (1:100, Cat# PA5-33850, Invitrogen, ThermoFisher Scientific, USA). Immunoreactivity was detected by using the Peroxidase Vectastain Elite ABC-HRP universal kit, according to the manufacturer's instructions. Cells were counterstained with hematoxylin and were visualized using light microscopy (Axiolab Karl Zeiss, Göttingen, Germany). HEK293 cells were used as negative control,²¹ while MDA-MB-231

and MCF-7 cells were employed as positive controls of H4R expression.^{12,14} The expression of H4R was further assessed in breast cancer cells by flow cytometry as previously described.¹² We used a primary rabbit anti-H4R antibody (1:100, Cat# ab97487; Abcam) followed by a secondary anti-rabbit antibody conjugated with FITC (1:80, Cat#F0382, Sigma Chemical Co., MO, USA). Samples were run on a BD Accuri C6 flow cytometer (BDB) and data was analyzed using the BD Accuri C6 software (BDB).

Histopathological and immunohistochemical analyses

Histopathological and immunohistochemical assessments were carried out on formalin-fixed paraffin-embedded tissue sections, which included representative samples of carcinomas and adjacent normal breast tissue. The diagnosis was established on hematoxylin and eosin sections by two board-certified pathologists separately. Histological grading and TNM staging were determined according to WHO classification.^{22,23} Tumors were categorized into low grade (grades 1 or 2) and high grade (grade 3), as previously described.²⁴

The blocks were cut in 5 µm sections and were immunolabeled with rabbit monoclonal antibodies directed against ER (Estrogen Receptor, clone SP1, 1:100, Cell Marque), PR (Progesterone Receptor, clone Y85, 1:30, Cell Marque), HER2 (Her2/Neu, clone SP3, 1:300, Cell Marque) and Ki67 (clone SP6, 1:200, Cell Marque), using an automated immunohistochemical staining equipment, according to the manufacturer's guidelines (Benchmark XT, Ventana, USA), and the standardized and approved procedure of the British Hospital Institution. Immunoreactivity was assessed blinded to

clinicopathological data, using a semiquantitative scoring system. The immunostaining scores for ER, PR and the algorithm for HER2 scoring were determined according to ASCO and CAP guidelines. Nuclear and membranous expression was considered positive for ER/PR and HER2, respectively. The threshold for the definition of TNBC was <1% immunopositivity of either ER or PR, and an immunoscore of 0 or 1+ for HER2 expression or 2+ in the absence of amplification by fluorescent in situ hybridization.^{4,22,24-26}

H4R immunostaining and scoring

The expression of H4R in tumors and peritumoral tissue was evaluated by immunohistochemical staining as it was previously described.¹⁴ Briefly, after deparaffinization the specimens were heated in a microwave in sodium citrate buffer (10 mM, pH 6.0) for antigen retrieval. After blocking, specimens were incubated with primary rabbit anti-H4R polyclonal antibody directed against the first cytoplasmic domain of human H4R (1:100, Cat# PA5-33850, Invitrogen) antibodies overnight in a humidified chamber at 4°C. Immunoreactivity was detected by using the Peroxidase Vectastain Elite ABC-HRP universal kit, according to the manufacturer's instructions. Pre-immune serum of the same animal species in which the secondary antibody was developed was used for blocking, and to replace the primary antibody to detect nonspecific binding of the secondary antibodies (PK-6200, Vector Laboratories, CA, USA). All specimens were processed following identical and standardized staining procedures.

The H4R immunoreactivity score was obtained by multiplying the intensity (negative, 0; weak, 1; moderate, 2; and strong, 3) by the percentage of stained

cells. H4R expression was considered to be “positive” if at least 5% of cell specimens showed membranous and/or granular cytoplasmic staining. All the evaluations were performed by consensus agreement of at least 2 specialized pathologists. Immunocompetent cells were considered internal positive controls in the specimens.¹¹ Visualization was performed with an optical microscope Leica ICC50 HD (Wetzlar, Germany). Photographs were taken at x100 and x400 magnification with Leica camera (Germany) and visualized with the Leica LAS EZ software (version 3.1.0, Leica Microsystem, Switzerland).

2.4. Statistical analysis

Statistical analyses were conducted using GraphPad Prism version 7.00 (San Diego, CA, USA). Mann-Whitney non-parametric test was used to compare average scores. Wilcoxon Matched-Pairs Signed Ranks Test was used for the statistical analysis of differences in protein expression between tumor-adjacent peritumoral normal tissue pairs. For determination of the association among different variables, Spearman’s rho correlation coefficients and two-tailed significance were determined. Log rank test and Gehan-Breslow-Wilcoxon test were performed for Kaplan-Meier survival. All statistical tests were two-sided, and a P-value <0.05 was significant.

RESULTS

H4R expression in human breast cancer samples

We have previously demonstrated the functional expression of H4R in TNBC experimental models in which H4R ligands showed anti-tumoral potential.¹²⁻¹⁷ However, the evidence of H4R expression and its role in human TNBC cancer progression has remained insufficient.

The potential clinical relevance of H4R in TNBC/basal-like tumors was assessed at a large scale by means of the genomic expression and clinical data obtained from publicly available datasets. Analyses of TCGA Pan-Cancer Atlas data set^{27,28} show that H4R mRNA expression was lower in the aggressive basal-like tumors compared with the more favorable clinical outcome luminal A (P=0.028) and normal breast-like tumors (P=0.018) (Fig. 1 A). Tumors were split into quartiles based on *H4R* expression and cancer subtypes, staging, and survival were investigated.

The highest quartile of *H4R* expression had a greater proportion of luminal A and normal breast-like tumors with lower proportion of basal-like compared to the lowest quartile (Fig. 1 B). Likewise, the evaluation of *H4R* gene alterations (including deletions, amplifications, and mutations) frequency in the different breast cancer subtypes obtained from cBioPortal web resource revealed that the vast proportion of gene alterations were observed in basal-like tumors (Supplementary Fig. 1 A). Interestingly, a significant reduced survival was observed in the group of breast cancer patients with H4R with at least one type

of gene alteration (Supplementary Fig. 1 A). Furthermore, higher levels of H4R mRNA expression were observed in early-stage breast cancer (Fig. 1 C).

These results suggest that the H4R seems to be particularly impaired in basal-like breast cancer. To deepen its role in tumor biology in this subtype and illustrate the potential prognostic value of H4R, survival rates based on progression or mortality were evaluated in basal-like breast cancer stratified by H4R low and high expression. Higher levels of H4R mRNA expression were significantly associated with improved progression free survival, and a non-significant increase in the overall survival (OS) in basal-like breast cancer (Fig. 1 D, E). Findings were confirmed using the Kaplan–Meier plotter database to evaluate the survival of basal-like cancer patients (probe 221170_x_at for H4R). The result indicated that a high level of H4R was significantly associated with improved relapse-free survival (HR 0.77, $P=0.016$, Fig. 1F) and OS (HR 0.64, $P = 0.019$, Fig. 1G) in basal-like cancer patients.

Although there is around an 80% overlap between triple negative and intrinsic basal-like breast cancer subtype, the basal-like classification is defined via gene expression analysis and to date is limited to the research setting. TNBC phenotype refers to the immunohistochemical classification of breast tumors lacking ER, PR, and HER2 protein expression and it is currently a reliable surrogate in the clinical setting.^{3,4,29}

To validate the transcriptomic data, we next evaluated the immunohistochemical protein expression of H4R in a small cohort of patients with TNBC. The specificity of the antibody was checked using HEK293 cells, which do not endogenously express H4R.²¹ As we have previously reported^{12,14}

we detected H4R in human MCF7 and MDA-MB-231 cells by immunocytochemistry, which served as positive controls (Fig. 1 H). Interestingly and in line with the results obtained of the expression of H4R in patient datasets, MCF-7 luminal-like breast cancer cells seemed to express higher levels of H4R compared to MDA-MB-231 TNBC cells. This upregulation of H4R in MCF-7 cells was confirmed by a semi-quantitative flow cytometric analysis (Fig. 1 I), and additionally investigating H4R mRNA expression (Cancer Cell Line Encyclopedia at cBioPortal) (Fig. 1 J).

Expression of H4R in TNBC and matched histologically normal breast tissue and its association with clinicopathological features

Next, we examined the relationship between H4R protein expression and its association with disease characteristics. Thirty TNBC specimens were analyzed in this study. Patient characteristics are summarized in Table 1. The clinicopathological features in the tumor samples were compared in the patient cohort. Negative nodal disease and unifocal TNBC were associated with a favorable prognosis (Supplementary Table 1).

H4R immunostaining shows a membranous and granular cytoplasmic pattern in the TNBC samples, which exhibited different levels of expression (Figure 2 A, Supplementary Fig. 2).

Twenty one of 30 tumors (70%) exhibited positive immunostaining for H4R with a score ranging between 5 and 180, while 9 tumors showed negative expression (Fig. 2 A-C). Additionally, the expression of H4R was analyzed in

the peritumoral normal tissue defined as the histologically normal tissue adjacent to the tumor. A positive H4R immunostaining was observed in 22 of 26 specimens (85%), exhibiting a score ranging from 5 to 270 (Fig. 2 B,C). Interestingly, there was a moderate positive correlation between the expression of H4R in the tumoral and peritumoral tissue (Fig. 2 C,D).

Considering that normal peritumoral tissue may exhibit alterations at molecular level that could be associated with cancer risk,³⁰⁻³² both types of samples were investigated. Elevated expression of H4R was demonstrated in relation to unifocal TNBC, which was significant in peritumoral histopathologically normal tissue (Table 2). No significant differences were detected between the H4R expression in tumor and peritumoral tissue and the histopathological grade, size, nodal status or the high proliferation index measured by Ki67 (Table 2). However, a negative correlation between H4R expression in peritumoral tissue and the number of lymph node involvement was found (Spearman r : -0.4793, P =0.015).

Survival studies showed that patients with H4R positivity have increased OS compared with H4R-negative specimens, which was significant considering H4R staining in peritumoral tissue (Fig. 2 E,F).

DISCUSSION

TNBC represents a major clinical therapeutic challenge. Recent data demonstrate the expression of H4R and its pathophysiological role in cancer, representing a potential molecular target for cancer therapeutics.^{11,13,14,17,33} This study provides evidence of the expression of H4R in TNBC and its potential association with prognosis.

TCGA is a publicly available database that shows the most important genomic changes in tumors of 33 types of cancers from thousands of patients, which notably contributes to accelerating our knowledge of the molecular basis of cancer with impacts in both cancer prevention and treatment. Using TCGA data, we have recently described the *H4R* gene expression in different types of tumors compared with matched-normal tissues. Depending on the cancer type, *H4R* seemed to be downregulated (e.g. colon adenocarcinoma, breast invasive carcinoma, bladder urothelial carcinoma), upregulated (e.g. hepatocellular, esophageal and kidney cancers) or unchanged (e.g. lung adenocarcinoma) compared to normal tissue.^{11,17,21,34-38}

In this study, we analyzed a large transcriptomic dataset associated with clinical features (TCGA Pan-Cancer Atlas) by means of cBioPortal for Cancer Genomics. The analysis of the H4R mRNA expression in different molecular subtypes of breast cancer demonstrated that H4R is downregulated in basal-like breast cancer compared with luminal A breast cancer and normal breast-like tumors, both favorable subtypes in terms of prognosis.^{3,29} In agreement with

these results, luminal MCF-7 cells showed higher H4R expression compared with basal-like MDA-MB-231 breast cancer cells.

An inverse relationship was evidenced when comparing the expression of *H4R* according to the neoplasm disease stage. A higher proportion of stage I non-spread breast cancer showed higher levels of *H4R* expression. In addition, the study of the alteration frequency of *H4R* gene in the breast cancer subtypes showed different percentages of alterations depending on the cancer subtype. The higher frequency of alterations, that include deletions and amplifications of the *H4R* gene, were observed in basal-like breast cancer compared to the other subtypes. Interestingly, survival analysis showed improved disease free survival in breast cancer patients without H4R gene alterations. Genomic alterations of this receptor in different cancer types have been described,¹⁷ however their role in carcinogenesis and in the response to therapeutics is completely unknown and deserves to be studied. These findings suggest that H4R may play a crucial role in breast cancer biology and progression, especially in the aggressive basal-like breast cancer. Therefore, Kaplan-Meier curves for basal-like breast cancer patients were stratified by H4R expression. The higher expression of H4R was associated with better survival clinical outcomes based on both progression and mortality events.

In line with these results, evidence from independent research groups demonstrated that potent H4R agonists reduced cell proliferation and events involved in the metastatic cascade in different cancer types.^{11,14,17,21,33,35,36,38} Therefore, H4R might contribute to improvements in cancer treatment in terms of a targeted therapy.

To corroborate the bioinformatic analyses, we investigated the protein expression of H4R in TNBC samples, according to pathology-based classification. To our knowledge, this study is the first to assess the immunohistochemical H4R expression specifically in human TNBC samples. Membranous and cytoplasmic H4R immunostaining was detected in 70% of TNBC samples. The expression of H4R was further demonstrated in the histologically normal breast tissue located adjacent to the carcinoma. The analysis of the expression of H4R in the peritumoral breast tissue revealed no significant differences with its expression in tumor epithelial cells, and a moderate positive correlation between the H4R score in the tumoral and peritumoral tissues.

Numerous reports suggest that histologically normal tissue adjacent to breast cancer may harbor molecular alterations, which could support tumorigenesis.^{30-32,39-45} In this connection, the identification in routine breast biopsies of a molecular marker in appearing normal tissue at risk for malignant transformation may have useful potential clinical application.^{12,32,44-45} H4R expression is inversely correlated with the number of regional lymph node metastases in peritumoral tissue. The number of involved axillary lymph nodes remains the dominant predictor of prognosis in breast cancer, overwhelming other factors and conditioning the decision of the adjuvant systemic treatment.⁴⁶⁻⁴⁹ Furthermore, multifocal TNBC was associated with reduced H4R expression in peritumoral tissue. Although a link between tumor focality and prognosis is not well understood, some studies show that multifocal lesions, when more than one tumor of the same origin arises in the same area of the breast, could be associated with a higher risk of recurrence.⁴⁶ In the absence of lymph node

metastases, tumor size and its histological grade, or proliferative index, contribute to sorting patients into groups according to cancer risk.⁴⁶ However, these parameters were neither prognostically important in terms of survival nor differentially modulated by H4R in our small patients' cohort.

Kaplan-Meier curves for OS of patients with TNBC were obtained according to the presence or absence of H4R in both tumor and peritumoral tissue. Patients with H4R expression had significantly better OS than those with undetectable levels of H4R just in peritumoral tissue. Presented data indicate that H4R expression in TNBC seems to be reduced or absent in more aggressive or disseminated tumors. We hypothesize that impairment of H4R expression in tumor-adjacent, histologically normal breast tissue could be present in breast epithelium as an early molecular change before clinical or pathological evidence of the neoplasm. Ongoing experimental studies are aimed at investigating the H4R expression in histologically normal tissue of breast cancer patients compared to normal epithelium of women without breast cancer as an approach to better understand the significance of H4R in carcinogenesis.

Our research has numerous limitations that should be described. First, the study was limited by a small sample size of a single institution. Some patients performed their treatments outside the institution or discontinued it, preventing follow-up data during a long period. Due to the small sample size, a meaningful statistical analysis of the correlation between H4R score and some clinicopathological parameters could not be possible. In addition, multivariate analysis is necessary to identify H4R as a potential independent predictor of clinical outcomes.

In conclusion, H4R transcriptomic data together with the immunohistochemical studies suggest that the H4R might represent a novel prognostic factor associated with aggressiveness and patient survival in TNBC, which could complement routine histopathological analysis. Furthermore, the detection of H4R in TNBC samples is clinically relevant considering that it could represent a promising therapeutic target for this aggressive and difficult-to-treat type of breast cancer. In this sense, this study serves as important data for the initiation of further studies to understand the significance of H4R in breast cancer biology and prognosis in large patient cohorts.

FIGURE LEGENDS

Figure 1. Bioinformatic analyses of the expression of *H4R* in breast cancer. mRNA expression levels of *H4R* were obtained from breast cancer datasets at the cBioportal for Cancer Genomics (TCGA PanCancer Atlas). **A)** *H4R* mRNA expression in different breast cancer (BRCA) subtypes. Box plots show the expression levels as log-transformed mRNA expression z-scores compared to the expression distribution of all samples (RNA Seq V2 RSEM). BRCA_Basal (n=171), BRCA_Her2 (n=78), BRCA_LumA (Luminal A, n=499), BRCA_LumB (Luminal B, n=197), BRCA_Normal (normal breast-like, n=36). a: P=0.028 vs. BRCA_Basal; b: P=0.018 vs. BRCA_Basal. Kruskal-Wallis test and Dunn's multiple comparisons test. **B)** Percentage of samples with different breast cancer subtypes based on *H4R* expression quartiles. Chi-squared test, P=0.047. **C)** Percentage of samples with different neoplasm disease stages American Joint Committee on Cancer (AJCC) code based on *H4R* expression quartiles. Chi-squared test, P=0.025. A (n=267): the lowest quartile, -2.02 to -1.14; B (n=268): -1.14 to -0.36; C (n=267): -0.35 to 0.45; D (n=268): the highest quartile, 0.45 to 6.06 (log RNA Seq V2 RSEM). **D, E)** Kaplan-Meier plots comparing the clinical outcomes of patients with high vs. low *H4R* expressing basal-like tumors (TCGA PanCancer Atlas). D) Progression free survival, and E) OS were evaluated for the lowest (A, n=42: -2.05 to -1.36 log RNA Seq V2 RSEM) and the highest (D, n=43: 0.17 to 4.18 log RNA Seq V2 RSEM) *H4R* expression quartiles. Mantel-Cox (Log-rank test). Progression free survival: P=0.045. OS: P=NS. **F, G)** Kaplan-Meier plots comparing the clinical outcomes of patients with high vs. low *H4R* expressing basal-like tumors (Kaplan–Meier Plotter). F) Relapse-free survival, and G) OS. Red line: patients with expression

levels above the median; black line: patients with expression levels below the median. Mantel-Cox (Log-rank test). Relapse free survival: P=0.016. OS: P=0.019. **H-J)** H4R expression in human cancer cell lines. H) Immunocytochemical detection of H4R in MDA-MB-231 and MCF-7 breast cancer cells. HEK293 cells were used as a negative control. x400 original magnification. Scale bar = 20 μ m. I) Immunofluorescence of H4R was evaluated by flow cytometry. Representative histograms are shown. J) H4R mRNA expression (RNA Seq RPKM) obtained at cBioPortal (Cancer Cell Line Encyclopedia, Broad 2019).⁵⁰

Figure 2. H4R expression in tumoral and peritumoral tissue of TNBC patients. **A)** Representative TNBC samples stained with hematoxylin & eosin (H&E) and H4R immunostaining. All corresponded to high grade invasive ductal carcinomas with variable amounts of ductal differentiation and numerous atypical mitoses. The score of H4R immunostaining was obtained by multiplying the intensity (negative, 0; weak, 1; moderate, 2; and strong, 3) by the percentage of stained cells. Negative expression of H4R (score 0), arrows indicate positive immunocompetent cells. Positive membranous and granular cytoplasmic staining of H4R with scores of 90 (3X30) and 180 (3X60) are shown. x100 and x400 original magnification. Scale bar = 20 μ m. Representative pictures of the scale of H4R intensities are shown in Supplementary Fig. 2. **B)** H4R immunostaining score in tumor (Score range: 5 - 180) and peritumoral breast tissue (Score range: 5 - 270). **C)** Similar H4R expression was seen in neoplastic cells (above) in comparison to normal ducts of the peritumoral breast lobules (below). H4R expression was always

membranous and cytoplasmic. x400 original magnification. **D)** Spearman's positive correlation between H4R expression in tumoral and peritumoral tissue. **E, F)** Kaplan-Meier survival curves according to the expression of H4R (Follow-up: 24 months). **E)** Log-rank and Mantel-Cox test: χ^2 (Chi-square)=0.001, P=NS, and the Gehan-Breslow-Wilcoxon test: $\chi^2=0.1656$, P = NS. **F)** Log-rank and Mantel-Cox test: $\chi^2 = 14.34$, P<0.001, and the Gehan-Breslow-Wilcoxon test: $\chi^2=12.78$, P<0.001.

Conflict of interest

The authors have no conflict of interest to declare.

Author Contributions

Conceived and designed the experiments: DS, JLU, VAM. Performed the experiments: DS, MTD, MBN, IO, FV, PD. Analyzed the data: DS, VAM. Contributed reagents/materials/analysis tools: JLU, AI, GE, VAM. Wrote the manuscript: DS, VAM. All authors approved the manuscript.

Funding

This work has been supported by grants from Fundación Alberto J. Roemmers, from National Agency for Scientific and Technological Promotion (PICT2018-03778, VAM), and the National Scientific and Technical Research Council (PIP-CONICET 11220200102459CO, VAM).

REFERENCES

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin.* 2021;71:209-249. <https://doi.org/10.3322/caac.21660>.
2. DeSantis CE, Ma J, Gaudet MM, Newman LA, Miller KD, Goding Sauer A, Jemal A, Siegel RL. Breast cancer statistics 2019. *CA Cancer J Clin.* 2019;69:438-451. <https://doi.org/10.3322/caac.21583>.
3. Prat A, Pineda E, Adamo B, Galván P, Fernández A, Gaba L, Díez M, Viladot M, Arance A, Muñoz M. Clinical implications of the intrinsic molecular subtypes of breast cancer. *Breast* 2015;24 Suppl 2:S26-35. <https://doi.org/10.1016/j.breast.2015.07.008>.
4. Goldhirsch A, Winer EP, Coates AS, Gelber RD, Piccart-Gebhart M, Thürlimann B, Senn HJ; Panel members. Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. *Ann Oncol.* 2013;24:2206-23. <https://doi.org/10.1093/annonc/mdt303>.
5. Bao C, Lu Y, Chen J, Chen D, Lou W, Ding B, Xu L, Fan W. Exploring specific prognostic biomarkers in triple-negative breast cancer. *Cell Death Dis.* 2019;10:807. <https://doi.org/10.1038/s41419-019-2043-x>.

6. Motamedi B, Rafiee-Pour HA, Khosravi MR, Kefayat A, Baradaran A, Amjadi E, Goli P. Prolactin receptor expression as a novel prognostic biomarker for triple negative breast cancer patients. *Ann Diagn Pathol.* 2020;46:151507. <https://doi.org/10.1016/j.anndiagpath.2020.151507>.
7. Pommier RM, Sanlaville A, Tonon L, Kielbassa J, Thomas E, Ferrari A, Sertier AS, Hollande F, Martinez P, Tissier A, Morel AP, Ouzounova M, Puisieux A. Comprehensive characterization of claudin-low breast tumors reflects the impact of the cell-of-origin on cancer evolution. *Nat Commun.* 2020;11:3431. <https://doi.org/10.1038/s41467-020-17249-7>.
8. Phillips L, Gill AJ, Baxter RC. Novel Prognostic Markers in Triple-Negative Breast Cancer Discovered by MALDI-Mass Spectrometry Imaging. *Front Oncol.* 2019;9:379. <https://doi.org/10.3389/fonc.2019.00379>.
9. Palleschi M, Maltoni R, Sarti S, Melegari E, Bravaccini S, Rocca A. Immunotherapy: The end of the "dark age" for metastatic triple-negative breast cancer?. *Breast J.* 2020;26:739-42. <https://doi.org/10.1111/tbj.13662>.
10. Medina VA, Rivera ES. Histamine receptors and cancer pharmacology. *Br J Pharmacol.* 2010;161:755-67. <https://doi.org/10.1111/j.1476-5381.2010.00961.x>.

11. Massari NA, Nicoud MB, Medina VA. Histamine receptors and cancer pharmacology: an update. *Br J Pharmacol.* 2020;177:516-38. <https://doi.org/10.1111/bph.14535>.
12. Medina V, Cricco G, Nuñez M, Martín G, Mohamad N, Correa-Fiz F, Sanchez-Jimenez F, Bergoc R, Rivera ES. Histamine-mediated signaling processes in human malignant mammary cells. *Cancer Biol Ther.* 2006;5:1462-71. <https://doi.org/10.4161/cbt.5.11.3273>.
13. Medina V, Croci M, Crescenti E, Mohamad N, Sanchez-Jiménez F, Massari N, et al. The role of histamine in human mammary carcinogenesis: H3 and H4 receptors as potential therapeutic targets for breast cancer treatment. *Cancer Biol Ther.* 2008;7:28-35. <https://doi.org/10.4161/cbt.7.1.5123>.
14. Martinel Lamas DJ, Croci M, Carabajal E, Crescenti EJ, Sambuco L, Massari NA, Bergoc RM, Rivera ES, Medina VA. Therapeutic potential of histamine H4 receptor agonists in triple-negative human breast cancer experimental model. *Br J Pharmacol.* 2013;170:188-99. <https://doi.org/10.1111/bph.12137>.
15. Nicoud MB, Sterle HA, Massari NA, Táquez Delgado MA, Formoso K, Herrero Ducloux MV, Martinel Lamas D, Cremaschi GA, Medina VA. Study of the antitumour effects and the modulation of immune response by histamine in breast cancer. *Br J Cancer.* 2020;122:348-60. <https://doi.org/10.1038/s41416-019-0636-x>.

16. Sterle HA, Nicoud MB, Massari NA, Táquez Delgado MA, Herrero Ducloux MV, Cremaschi GA, Medina VA. Immunomodulatory role of histamine H4 receptor in breast cancer. *Br J Cancer*. 2019;120:128-38. <https://doi.org/10.1038/s41416-018-0173-z>.
17. Nicoud MB, Formoso K, Medina VA. Pathophysiological Role of Histamine H4 Receptor in Cancer: Therapeutic Implications. *Front Pharmacol*. 2019;10:556. <https://doi.org/10.3389/fphar.2019.00556>.
18. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, Antipin Y, Reva B, Goldberg AP, Sander C, Schultz N. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012;2:401-4. <https://doi.org/10.1158/2159-8290.CD-12-0095>.
19. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, Cerami E, Sander C, Schultz N. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal*. 2013;6:pl1. <https://doi.org/10.1126/scisignal.2004088>.
20. Györfy B. Survival analysis across the entire transcriptome identifies biomarkers with the highest prognostic power in breast cancer. *Comput Struct Biotechnol J*. 2021;19:4101-4109. <https://doi.org/10.1016/j.csbj.2021.07.014>.

21. Massari NA, Nicoud MB, Sambuco L, Cricco GP, Martinel Lamas DJ, Herrero Ducloux MV, Blanco H, Rivera ES, Medina VA. Histamine therapeutic efficacy in metastatic melanoma: Role of histamine H4 receptor agonists and opportunity for combination with radiation. *Oncotarget*. 2017;8:26471-91. <https://doi.org/10.18632/oncotarget.15594>.

22. Rakha EA, Allison KH, Ellis IO, Horii B, Masuda S, Penault-Llorca F, Tsuda H, Vincent Salomon A. Invasive breast carcinoma: General overview. In: WHO Classification of Tumours Editorial Board, editors. *WHO Classification of Tumours, Breast Tumours*, 5th edition. Lyon: IARC Press; 2019, p. 82-100.

23. Union for international cancer control (UICC). Breast Tumours. In: Brierley J, Gospodarowicz MK, Wittekind Ch, editors. *TNM Classification of malignant tumours*, 8th edition. Oxford: Wiley & sons; 2017, p.151-7.

24. Hammond ME, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, Fitzgibbons PL, Francis G, Goldstein NS, Hayes M, Hicks DG, Lester S, Love R, Mangu PB, McShane L, Miller K, Osborne CK, Paik S, Perlmutter J, Rhodes A, Sasano H, Schwartz JN, Sweep FC, Taube S, Torlakovic EE, Valenstein P, Viale G, Visscher D, Wheeler T, Williams RB, Wittliff JL, Wolff AC. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *Arch Pathol Lab Med*. 2010;134:907-22. <https://doi.org/10.5858/134.6.907>.

25. Ilie SM, Bacinschi XE, Botnariuc I, Anghel RM. Potential clinically useful prognostic biomarkers in triple-negative breast cancer: preliminary results of a retrospective analysis. *Breast Cancer* (Dove Med Press). 2018;10:177-94.

<https://doi.org/10.2147/BCTT.S175556>.

26. Fitzgibbons PL, Connolly JL. Template for Reporting Results of Biomarker Testing of Specimens From Patients With Carcinoma of the Breast. College of American Pathologists (CAP). Version Breast Biomarkers 1.4.1.1. available in: <https://www.cap.org/protocols-and-guidelines/cancer-reporting-tools/cancer-protocol-templates>; 2021.

27. Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, Shen R, Taylor AM, Cherniack AD, Thorsson V, Akbani R, Bowlby R, Wong CK, Wiznerowicz M, Sanchez-Vega F, Robertson AG, Schneider BG, Lawrence MS, Noushmehr H, Malta TM. Cancer Genome Atlas Network, Stuart JM, Benz CC, Laird PW. Cell-of-Origin Patterns Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. *Cell*. 2018;173:291-304.

<https://doi.org/10.1016/j.cell.2018.03.022>.

28. Liu J, Lichtenberg T, Hoadley KA, Poisson LM, Lazar AJ, Cherniack AD, Kovatich AJ, Benz CC, Levine DA, Lee AV, Omberg L, Wolf DM, Shriver CD, Thorsson V; Cancer Genome Atlas Research Network, Hu H. An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. *Cell*. 2018;173:400-416.

<https://doi.org/10.1016/j.cell.2018.02.052>.

29. Yersal O, Barutca S. Biological subtypes of breast cancer: Prognostic and therapeutic implications. *World J Clin Oncol.* 2014;5:412-24. <https://doi.org/10.5306/wjco.v5.i3.412>.
30. Chen DT, Nasir A, Culhane A, Venkataramu C, Fulp W, Rubio R, Wang T, Agrawal D, McCarthy SM, Gruidl M, Bloom G, Anderson T, White J, Quackenbush J, Yeatman T. Proliferative genes dominate malignancy-risk gene signature in histologically-normal breast tissue. *Breast Cancer Res Treat.* 2010;119:335-46. <https://doi.org/10.1007/s10549-009-0344-y>.
31. Graham K, de las Morenas A, Tripathi A, King C, Kavanah M, Mendez J, Stone M, Slama J, Miller M, Antoine G, Willers H, Sebastiani P, Rosenberg CL. Gene expression in histologically normal epithelium from breast cancer patients and from cancer-free prophylactic mastectomy patients shares a similar profile. *Br J Cancer.* 2010;102:1284-93. <https://doi.org/10.1038/sj.bjc.6605576>.
32. Tripathi A, King C, de la Morenas A, Perry VK, Burke B, Antoine GA, Hirsch EF, Kavanah M, Mendez J, Stone M, Gerry NP, Lenburg ME, Rosenberg CL. Gene expression abnormalities in histologically normal breast epithelium of breast cancer patients. *Int J Cancer.* 2008;122:1557-66. <https://doi.org/10.1002/ijc.23267>.
33. Medina VA, Brenzoni PG, Lamas DJ, Massari N, Mondillo C, Nunez MA, Pignataro O, Rivera ES. Role of histamine H4 receptor in breast cancer cell

proliferation. *Front Biosci (Elite Ed)*. 2011;3:1042-60.
<https://doi.org/10.2741/e310>.

34. Boer K, Helinger E, Helinger A, Pocza P, Pos Z, Demeter P, Baranyai Z, Dede K, Darvas Z, Falus A. Decreased expression of histamine H1 and H4 receptors suggests disturbance of local regulation in human colorectal tumours by histamine. *Eur J Cell Biol*. 2008;87:227-36.
<https://doi.org/10.1016/j.ejcb.2007.12.003>.

35. Fang Z, Yao W, Xiong Y, Li J, Liu L, Shi L, Zhang W, Zhang C, Nie L, Wan J. Attenuated expression of HRH4 in colorectal carcinomas: a potential influence on tumor growth and progression. *BMC Cancer*. 2011;11:195.
<https://doi.org/10.1186/1471-2407-11-195>.

36. Meng F, Han Y, Staloch D, Francis T, Stokes A, Francis H. The H4 histamine receptor agonist, clobenpropit, suppresses human cholangiocarcinoma progression by disruption of epithelial mesenchymal transition and tumor metastasis. *Hepatology*. 2011;54:1718-28.
<https://doi.org/10.1002/hep.24573>.

37. Zhang C, Xiong Y, Li J, Yang Y, Liu L, Wang W, , Wang L, Li M, Fang Z. Deletion and down-regulation of HRH4 gene in gastric carcinomas: a potential correlation with tumor progression. *PLoS One*. 2012;7:e31207.
<https://doi.org/10.1371/journal.pone.0031207>.

38. He GH, Ding JQ, Zhang X, Xu WM, Lin XQ, Huang MJ, Feng J, Wang P, Cai WK. Activation of histamine H4 receptor suppresses the proliferation and invasion of esophageal squamous cell carcinoma via both metabolism and non-metabolism signaling pathways. *J Mol Med (Berl)*. 2018;96:951-64. <https://doi.org/10.1007/s00109-018-1676-z>.

39. Lewis CM, Cler LR, Bu DW, Zöchbauer-Müller S, Milchgrub S, Naftalis EZ, , Leitch AM, Minna JD, Euhus DM. Promoter hypermethylation in benign breast epithelium in relation to predicted breast cancer risk. *Clin Cancer Res*. 2005;11:166-72.

40. Deng G, Lu Y, Zlotnikov G, Thor AD, Smith HS. Loss of heterozygosity in normal tissue adjacent to breast carcinomas. *Science*. 1996;274:2057-9. <https://doi.org/10.1126/science.274.5295.2057>.

41. Botti C, Pescatore B, Mottolese M, Sciarretta F, Greco C, Di Filippo F, Gandolfo GM, Cavaliere F, Bovani R, Varanese A, Cianciulli AM. Incidence of chromosomes 1 and 17 aneusomy in breast cancer and adjacent tissue: an interphase cytogenetic study. *J Am Coll Surg*. 2000;190:530-9. [https://doi.org/10.1016/s1072-7515\(00\)00252-0](https://doi.org/10.1016/s1072-7515(00)00252-0).

42. Larson PS, de las Morenas A, Bennett SR, Cupples LA, Rosenberg CL. Loss of heterozygosity or allele imbalance in histologically normal breast epithelium is distinct from loss of heterozygosity or allele imbalance in co-

existing carcinomas. *Am J Pathol.* 2002;161:283-90.
[https://doi.org/10.1016/S0002-9440\(10\)64180-6](https://doi.org/10.1016/S0002-9440(10)64180-6).

43. Li Z, Moore DH, Meng ZH, Ljung BM, Gray JW, Dairkee SH. Increased risk of local recurrence is associated with allelic loss in normal lobules of breast cancer patients. *Cancer Res.* 2002;62:1000-3.

44. Trujillo KA, Heaphy CM, Mai M, Vargas KM, Jones AC, Vo P, Butler KS, Joste NE, Bisoffi M, Griffith JK. Markers of fibrosis and epithelial to mesenchymal transition demonstrate field cancerization in histologically normal tissue adjacent to breast tumors. *Int J Cancer.* 2011;129:1310-21.
<https://doi.org/10.1002/ijc.25788>.

45. Aran D, Camarda R, Odegaard J, Paik H, Oskotsky B, Krings G, , Goga A, Sirota M, Butte AJ. Comprehensive analysis of normal adjacent to tumor transcriptomes. *Nat Commun.* 2017;8:1077. <https://doi.org/10.1038/s41467-017-01027-z>.

46. Arciero CA and Styblo TM. Clinically Established Prognostic Factors in Breast Cancer, Chapter 18, Section VI. In: Bland KI, Copeland EM, Klimberg VS, Gradishar WJ, editors. *The Breast: Comprehensive management of benign and malignant diseases*, 5th edition. Philadelphia: Elsevier; 2018, p.250-8.

47. Henderson IC, Patek AJ. The relationship between prognostic and predictive factors in the management of breast cancer. *Breast Cancer Res Treat.* 1998;52:261-88. <https://doi.org/10.1023/a:1006141703224>.

48. Carter CL, Allen C, Henson DE. Relation of tumor size, lymph node status, and survival in 24,740 breast cancer cases. *Cancer.* 1989;63:181-7. <https://doi.org/10.1136/bmj.300.6721.346>.


49. Fisher B, Bauer M, Wickerham DL, Redmond CK, Fisher ER, Cruz AB, Foster R, Gardner B, Lerner H, Margolese R, Poisson R, Shibata H, Volk H. Relation of number of positive axillary nodes to the prognosis of patients with primary breast cancer. An NSABP update. *Cancer.* 1983;52:1551-7. [https://doi.org/10.1002/1097-0142\(19831101\)52:9<1551::aid-cncr2820520902>3.0.co;2-3](https://doi.org/10.1002/1097-0142(19831101)52:9<1551::aid-cncr2820520902>3.0.co;2-3).

50. Ghandi M, Huang FW, Jané-Valbuena J, Kryukov GV, Lo CC, McDonald ER 3rd, Barretina J, Gelfand ET, Bielski CM, Li H, Hu K, Andreev-Drakhlin AY, Kim J, Hess JM, Haas BJ, Aguet F, Weir BA, Rothberg MV, Paoletta BR, Lawrence MS, Akbani R, Lu Y, Tiv HL, Gokhale PC, de Weck A, Mansour AA, Oh C, Shih J, Hadi K, Rosen Y, Bistline J, Venkatesan K, Reddy A, Sonkin D, Liu M, Lehar J, Korn JM, Porter DA, Jones MD, Golji J, Caponigro G, Taylor JE, Dunning CM, Creech AL, Warren AC, McFarland JM, Zamanighomi M, Kauffmann A, Stransky N, Imielinski M, Maruvka YE, Cherniack AD, Tsherniak A, Vazquez F, Jaffe JD, Lane AA, Weinstock DM, Johannessen CM, Morrissey MP, Stegmeier F, Schlegel R, Hahn WC, Getz G, Mills GB, Boehm JS, Golub TR, Garraway

LA, Sellers WR. Next-generation characterization of the Cancer Cell Line Encyclopedia. Nature. 2019;569:503-508. <https://doi.org/10.1038/s41586-019-1186-3>.


Figure 1. Bioinformatic analyses of the expression of H4R in breast cancer. mRNA expression levels of H4R were obtained from breast cancer datasets at the cBioportal for Cancer Genomics (TCGA PanCancer Atlas). A) H4R mRNA expression in different breast cancer (BRCA) subtypes. Box plots show the expression levels as log-transformed mRNA expression z-scores compared to the expression distribution of all samples (RNA Seq V2 RSEM). BRCA_Basal (n=171), BRCA_Her2 (n=78), BRCA_LumA (Luminal A, n=499), BRCA_LumB (Luminal B, n=197), BRCA_Normal (normal breast-like, n=36). a: P=0.028 vs. BRCA_Basal; b: P=0.018 vs. BRCA_Basal. Kruskal-Wallis test and Dunn's multiple comparisons test. B) Percentage of samples with different breast cancer subtypes based on H4R expression quartiles. Chi-squared test, P=0.047. C) Percentage of samples with different neoplasm disease stages American Joint Committee on Cancer (AJCC) code based on H4R expression quartiles. Chi-squared test, P=0.025. A (n=267): the lowest quartile, -2.02 to -1.14; B (n=268): -1.14 to -0.36; C (n=267): -0.35 to 0.45; D (n=268): the highest quartile, 0.45 to 6.06 (log RNA Seq V2 RSEM). D, E) Kaplan-Meier plots comparing the clinical outcomes of patients with high vs. low H4R expressing basal-like tumors (TCGA PanCancer Atlas). D) Progression free survival, and E) OS were evaluated for the lowest (A, n=42: -2.05 to -1.36 log RNA Seq V2 RSEM) and the highest (D, n=43: 0.17 to 4.18 log RNA Seq V2 RSEM) H4R expression quartiles. Mantel-Cox (Log-rank test). Progression free survival: P=0.045. OS: P=NS. F, G) Kaplan-Meier plots comparing the clinical outcomes of patients with high vs. low H4R expressing basal-like tumors (Kaplan-Meier Plotter). F) Relapse-free survival, and G) OS. Red line: patients with expression levels above the median; black line: patients with expression levels below the median. Mantel-Cox (Log-rank test). Relapse free survival: P=0.016. OS: P=0.019. H-J) H4R expression in human cancer cell lines. H) Immunocytochemical detection of H4R in MDA-MB-231 and MCF-7 breast cancer cells. HEK293 cells were used as a negative control. x400 original magnification. Scale bar = 20 μ m. I) Immunofluorescence of H4R was evaluated by flow cytometry. Representative histograms are shown. J) H4R mRNA expression (RNA Seq RPKM) obtained at cBioPortal (Cancer Cell Line Encyclopedia, Broad 2019).50

Figure 1




Supplementary Figure 1

A



B



Supplementary Fig. 1. *H4R* genomic alterations in breast cancer. A) *H4R* alteration event frequency in breast cancer subtypes. The alterations types include deletion, amplification, multiple alteration, and mutation. Data was obtained from cBioPortal web resource (TCGA PanCancer Atlas). BRCA_Basal (6 of 171), BRCA_Her2 (0 of 78), BRCA_LumA (Luminal A, 1 of 499), BRCA_LumB (Luminal B, 3 of 197), BRCA_Normal (normal breast-like, 0 of 36). $P=0.004$. **B)** Kaplan-Meier plot comparing the clinical outcome of patients with altered vs. unaltered *H4R* gene (TCGA PanCancer Atlas). Mantel-Cox (Logrank test). Disease free survival: $P=0.047$.

Table 1: Clinicopathological characteristics of the TNBC patients

| Population | Variables | Patient Number N=30 | Proportion (%) |
|------------------------------|---------------------------|--------------------------------|---------------------------|
| Clinical features | | | |
| Age (years) | Mean/Range | 52.3 (25-69) | |
| Tumor laterality | Right Breast | 12 | 40 |
| | Left Breast | 18 | 60 |
| Tumor Focality | Unifocal | 21 | 70 |
| | Multifocal | 9 | 30 |
| Type of surgery | Breast conserving surgery | 23 | 76.6 |
| | Mastectomy | 7 | 23.3 |
| Pathological features | | | |
| Size (cm) | Mean/Range | 2.06 (0.4- 4.5) | |
| Histopathology | Invasive ductal carcinoma | 26 | 86.6 |
| | Other type | 4 | 13.3 |
| Histologic grade | High grade | 21 | 70 |
| | Low grade | 9 | 30 |
| Lymphovascular invasion | No | 21 | 70 |
| | Yes | 9 | 30 |
| Accompanying In Situ pattern | No | 8 | 26.6 |
| | Yes | 22 | 73.3 |
| Lymph node metastases | No | 18 | 60 |
| | Yes | 12 | 40 |
| Ki67 | ≤20% | 6 | 20 |
| | >20% | 24 | 80 |
| Histologic stage | I-II | 25 | 83.3 |
| | III | 5 | 16.6 |
| Recurrence | No | 24 | 80 |
| | Yes | 6 | 20 |

Figure 2. H4R expression in tumoral and peritumoral tissue of TNBC patients. A) Representative TNBC samples stained with hematoxylin & eosin (H&E) and H4R immunostaining. All corresponded to high grade invasive ductal carcinomas with variable amounts of ductal differentiation and numerous atypical mitoses. The score of H4R immunostaining was obtained by multiplying the intensity (negative, 0; weak, 1; moderate, 2; and strong, 3) by the percentage of stained cells. Negative expression of H4R (score 0), arrows indicate positive immunocompetent cells. Positive membranous and granular cytoplasmic staining of H4R with scores of 90 (3X30) and 180 (3X60) are shown. x100 and x400 original magnification. Scale bar = 20 μ m. Representative pictures of the scale of H4R intensities are shown in Supplementary Fig. 2. B) H4R immunostaining score in tumor (Score range: 5 - 180) and peritumoral breast tissue (Score range: 5 - 270). C) Similar H4R expression was seen in neoplastic cells (above) in comparison to normal ducts of the peritumoral breast lobules (below). H4R expression was always membranous and cytoplasmic. x400 original magnification. D) Spearman's positive correlation between H4R expression in tumoral and peritumoral tissue. E, F) Kaplan-Meier survival curves according to the expression of H4R (Follow-up: 24 months). E) Log-rank and Mantel-Cox test: χ^2 (Chi-square)=0.001, P=NS, and the Gehan-Breslow-Wilcoxon test: $\chi^2=0.1656$, P = NS. F) Log-rank and Mantel-Cox test: $\chi^2 = 14.34$, P<0.001, and the Gehan-Breslow-Wilcoxon test: $\chi^2=12.78$, P<0.001.


Figure 2

H&E

H4R (x100)

H4R (x400)

A




(score 0)

(Score 90)


(Score 180)

H4R expression


B



C




D



E

**Overall Survival
H4R tumor expression**



F

**Overall Survival
H4R peritumor expression**





Table 2: H4R expression according to different clinicopathological parameters in TNBC patients.

| Clinicopathological parameter | | H4R expression | | | |
|-------------------------------|------------|----------------|-----------------|-------------------|-----------------|
| | | Tumor Score | | Peritumoral Score | |
| | | Median (IQR) | <i>P</i> -value | Median (IQR) | <i>P</i> -value |
| Tumor Focality | Unifocal | 50 (12.5-95) | NS | 60 (7.5-160) | 0.011 |
| | Multifocal | 0 (0-80) | | 5 (0-17.5) | |
| Histologic grade | High | 40 (0-82.5) | NS | 10 (5-125) | NS |
| | Low | 35 (0-142.5) | | 17.5 (1-140) | |
| LN metastases | No | 50 (3.7-105) | NS | 80 (5-160) | NS |
| | Yes | 25 (0-72.5) | | 10 (1-48.7) | |
| Ki67 | ≤20% | 50 (15-110) | NS | 140 (62.5-170) | NS |
| | >20% | 40 (0-87.5) | | 10 (5-55) | |
| Tumor size | >2 cm | 35 (0-115) | NS | 10 (5-35) | NS |
| | ≤2 cm | 45 (0-80) | | 100 (5-160) | |

LN: Lymph node, NS: not significant, IQR: interquartile range.

Mann Whitney's Test.

Supplementary Figure 2



Supplementary Fig. 2. Representative pictures of the scale of H4R intensities. Evaluations were performed in consensus agreement by at least two experienced pathologists. H4R immunostaining intensity: negative, 0; weak, 1; moderate, 2; and strong, 3.

Supplementary Table 1: Overall survival according to different clinicopathological parameters in TNBC patients.

| Clinicopathological parameter | | Survival | | | | |
|-------------------------------|------------|----------|---------------|-----------------|---------------|-----------------|
| | | Effect | Log-rank test | | Wilcoxon test | |
| | | | χ^2 | <i>P</i> -value | χ^2 | <i>P</i> -value |
| Tumor Focality | Unifocal | ↑ | 3.85 | 0.0497 | 3.67 | 0.0555 |
| | Multifocal | | | | | |
| Histologic grade | High | | 0.61 | NS | 0.66 | NS |
| | Low | | | | | |
| LN metastases | No | ↑ | 4.04 | 0.0444 | 6.92 | 0.0085 |
| | Yes | | | | | |
| Ki67 | ≤20% | | 1.01 | NS | 0.98 | NS |
| | >20% | | | | | |
| Tumor size | >2 cm | | 0.001 | NS | 0 | NS |
| | ≤2 cm | | | | | |

LN: Lymph node, NS: not significant