

Vigo, Daniel E., ; Simonelli, Guido ; Tuñón, Ianina ; Pérez Chada, Daniel ; Cardinali, Daniel P. ; Golombek, Diego

School characteristics, child work, and other daily activities as sleep deficit predictors in adolescents from households with unsatisfied basic needs

Preprint del documento publicado en
Mind Brain and Education Vol.8 N° 4, 2014

Este documento está disponible en la Biblioteca Digital de la Universidad Católica Argentina, repositorio institucional desarrollado por la Biblioteca Central "San Benito Abad". Su objetivo es difundir y preservar la producción intelectual de la Institución.

La Biblioteca posee la autorización de los autores y de la editorial para su divulgación en línea.

Cómo citar el documento:

Vigo, DE, Simonelli, G, Tuñón, I. et al. School characteristics, child work, and other daily activities as sleep deficit predictors in adolescents from households with unsatisfied basic needs [en línea]. Preprint del documento publicado en Mind brain and education 2014;8(4). Disponible en :<http://bibliotecadigital.uca.edu.ar/repositorio/investigacion/school-characteristics-child-work.pdf> [Fecha de consulta:]

School characteristics, child work, and other daily activities as sleep deficit predictors in adolescents from households with unsatisfied basic needs

Daniel E. Vigo (1,5), Guido Simonelli (1), Ianina Tuñón (2), Daniel Pérez Chada (3), Daniel P. Cardinali (1,5), Diego Golombek (4,5).

(1) Laboratorio de Neurociencia Aplicada, Departamento de Docencia e Investigación, Facultad de Ciencias Médicas, Pontificia Universidad Católica Argentina.

(2) Observatorio de la Deuda Social Argentina, Pontificia Universidad Católica Argentina.

(3) Departamento de Neumonología, Hospital Universitario Austral.

(4) Laboratorio de Cronobiología, Universidad Nacional de Quilmes.

(5) Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

Corresponding author

Daniel E. Vigo

Facultad de Ciencias Médicas, Pontificia Universidad Católica Argentina

Alicia Moreau de Justo 1500

C1107AAZ Ciudad Autónoma de Buenos Aires

Argentina.

Tel. +54 11 43490200 ext 1152; fax: +54 11 43490200.

E-mail address: dvigo@conicet.gov.ar

Key words

Sleep, adolescents, school, poverty, child labour

Abstract

Sleep in adolescents has been shown to be an important factor when looking at physical, mental and social well-being. Little evidence is found regarding sleep patterns in adolescents from households facing extreme poverty, where conditions like crowding, poor housing, sanitation or education, and precarious employment set an adverse environment for sleep. In this study we sought to assess in a Argentine, nation-wide sample comprising 1682 adolescents, how the presence of extreme poverty - as defined by the presence of unsatisfied basic needs (UBN) - affects the relationship between sleep duration and school, work and other daily activities. A global high prevalence of short sleeping time, a slight increase of sleep time in adolescents with UBN, and different patterns of wake activities that predict sleep deficit, depending on the presence of UBN, were found. The poor academic achievement, increased risk of accidents and adverse health outcomes associated with sleep deprivation, support the view that sleep is an additional unsatisfied basic need that worsens living conditions at this age. The results may help to design public health policies that contribute to ameliorate this adverse situation.

Introduction

Sleep in adolescents has been shown to be an important factor when looking at physical, mental and social well-being. Adolescents with short sleep duration are at an increased risk of accidental injuries (Carskadon, Acebo, & Jenni, 2004). Reduced sleep is considered a risk factor for overweight and obesity, as well as for type 2 diabetes and cardiovascular disease (Cappuccio et al., 2008; Perez-Chada, Drake, Perez-Lloret, Videla, & Cardinali, 2009). A number of studies have demonstrated that sleep of insufficient duration or quality is associated with different negative mental health outcomes that include anxiety, depression, and bipolar disorders (Mindell & Meltzer, 2008). Youths who experienced sleep problems had greater odds of interpersonal dysfunction (Roberts, Roberts, & Chen, 2001). Also, late bed and rise times, erratic sleep-wake schedules, shortened total sleep time, and poor sleep quality are negatively associated with academic performance (Wolfson & Carskadon, 2003; Perez-Chada et al., 2007; Perez-Lloret et al., 2013).

Several factors affect sleep in this age group. Developmental changes due to intrinsic regulatory mechanisms in the homeostatic and circadian sleep processes delay the timing of sleep. Psychosocial issues as self-selected bedtimes, academic pressure, the use of technological resources and social networking in the evening also delay bed time. On the other side, social pressures determine an early rise time for starting school day timely. Therefore, adolescents are asked to be awake at an inadequate circadian phase and sleep too little (Carskadon, 2011).

In addition, it was observed that a “sleep disparity” exists in the general population since poor sleep is strongly associated with poverty and race. A differential vulnerability to factors such as health, education and employment was suggested as mediator for this effect (Patel, Grandner, Xie, Branas, & Gooneratne, 2010; Cardinali et al., 2013). Adolescents from lower socio-economic households have poorer and less consistent sleep than those from more favored backgrounds, being the neighborhood and home conditions the aspects associated with these observations (Marco, Wolfson, Sparling, & Azuaje, 2011). This is aggravated by the fact that in poorer households child work is much more common, which tends to increase the risk for sleep problems since adolescents are involved in dual duties that further limit their time for sleep (Teixeira, Fischer, & Lowden, 2006; Vinha D, Cavalcante JA, & Andrade M, 2010; Fischer, Nagai, & Teixeira, 2008). According to the International Labour Organization (ILO) around 168 million children aged 5-17 years worldwide are engaged in child labour, accounting

for almost 11% of the child population as a whole (International Labour Office - International Programme on the Elimination of Child Labour (IPEC), 2013). In Argentina, approximately 20% of teens between 13 and 17 years was involved in economical activities and 14% was involved in intensive domestic work in 2012 (Tuñón, 2013).

In this context, less evidence was found regarding sleep problems in adolescents from households facing extreme poverty conditions, where conditions like crowding, poor housing, sanitation or educational level, and a precarious employment set an adverse environment for sleep. For instance, slum dwellers report bad indexes of sleep quality, sleep latency, sleep duration, sleep efficiency, sleep disturbances, daytime dysfunction that improve after minimal changes in the quality of basic housing. Understanding sleep patterns in this population could help to define interventions to reduce sleep disparity (Simonelli et al., 2013). Therefore, we sought to assess in a nation-wide sample, how the presence of extreme poverty - as defined by the presence of unsatisfied basic needs (UBN) - affects the already known relationship of sleep duration with school and working schedules, controlling for the practice of other daily activities.

Methods

Subjects and design

The analysis employ data from the 2012 Argentine Social Debt Barometer (Barómetro de la Deuda Social Argentina), Pontificia Universidad Católica Argentina, a nationwide urban probability sample of adults residing Argentina aged 18 and over. Participants were selected by multistage cluster random sampling based on urban agglomeration, housing conditions and socioeconomic status. In the first stage demographic criteria (by geographic region and size) was used to select 20 urban agglomerations of 80,000 inhabitants or more. Second, stratified random sampling was used in conjunction with a variable-radius plot, with probability proportional to the size of the 18 year old and older population. At the third stage, random systematic sampling was used to select each house inside the radius plot. A trained interviewer visited the houses, and individuals invited to participate on each house were randomly selected through a quota system of age and sex. Finally, 5636 individuals (46.4% males and 53.6% females) gave oral informed consent and participated in the survey. The information regarding the teenagers that lived in the house was gathered only when the respondent was the parent

or legal guardian of them. For the analysis, the cases were weighted to reflect each subgroup's actual proportion in the population.

Measures

Demographics. The Unsatisfied Basic Needs (UBN) index was used as an indicator of extreme poverty. The presence of UBN was defined when children or adolescents lived in a household that presented at least one of the following: a. more than three people per room (crowding); b. house built with irregular materials or living in a rented room (housing); c. not having an indoor flush toilet (sanitation); d. having a child between 6-12 years old who is not attending to school (school attendance); and e. households with four or more people per employed person and the maximum educational level of the household head being elementary. The sample was divided by the presence of UBN. Numerical age and percentage of female gender was reported.

Sleep. Questions regarding sleep included bedtime (lights off), risetime, and total amount of sleep during the day (nap duration). The variables derived from these questions were night sleep (risetime – bedtime), nap habit (nap duration > 0), total sleep time (TST = night sleep + nap duration), and sleep deficit (TST < 9hs for adolescents).

School. Questions regarding school characteristics included school starting time (morning, afternoon, or night), school distance (in city blocks) and school day length (simple or double shift). School distance > 1km was derived from the variable school distance.

Child labour. The propensity of being engaged in intensive domestic work (“home work”) was defined when the adolescent usually has the responsibility of: a. house attendance (clean and dust, wash and iron, etc.); b. cooking; c. taking care of siblings; or d. undertaking shopping, running errands, or fetching wood or water. The propensity of being engaged in economical activities (“paid work”) was considered when the child or adolescent helped his/her parents in their job, did an activity by him/herself for earning money, or had a job as an employee or trainee, within the last 12 months.

Other activities. Activities that may interfere with sleep were also reported, including a. the daily use of screens (TV, computers, play station) for more than two hours (“screen > 2hs);

b. daily outdoor playing (biking, skating, running, climbing, football, etc.) for more than two hours (“outdoor > 2hs”); c. non-scholar intellectual activities (music, painting, dance, foreign languages, chess, computer courses, scholar support courses, etc.) within the last 30 days (“extra-curricular intellectual activities”); and d. non-scholar sport or physical activities (football, basket, bike, swimming, skate, etc.) within the last 30 days (“extra-curricular physical activities”).

Statistical analyses

Numerical variables were reported as mean \pm standard error and categorical variables were reported as frequency (percentage). Differences between adolescents with or without UBN and between adolescents with or without sleep deficit were assessed through independent T-tests (numerical variables) or chi-square tests (categorical variables). A binary logistic regression model was conducted to determine independent predictors of sleep deficit in subjects with or without UBN, introducing independent factors by means of a stepwise forward conditional method.

Results

The sample included 1682 teenagers between 13 and 17 years old. Table 1 shows demographics, sleep duration, school characteristics and other daily activities of the sample. The presence of sleep deficit was observed in 38% of the whole sample, with a mean bedtime at 23.4 ± 0.04 hours and a mean risetime at 6.8 ± 0.03 hours, comprising a mean night sleep time of 8.4 ± 0.03 hours. Nap habit was present in 15.2% of these subjects, determining a mean total sleep time of 7.4 ± 0.03 hours. Most adolescents assist to morning school. Evening students were very few, so they were excluded from further statistical analyses. The percentage of teens who were involved in work was 13% (home) and 18% (rented work). Screen use > 2hs was the activity that involved the highest percentage of the studied sample (Table 1).

A different pattern of sleep-wake characteristics associated with sleep deficit is observed depending on the presence of UBN (Table 1). UBN was significantly associated with a lower percentage of sleep deficit ($X^2 = 5.35$, $p = .021$), mild later rise-time ($t = -3.40$, $p = .001$),

increased night sleep ($t = -4.40$, $p < .001$) and total sleep time ($t = -4.11$, $p < .001$), lower percentage of adolescents that assist to distant schools ($X^2 = 23.1$, $p < .001$), higher percentage of adolescents that work at home ($X^2 = 73.0$, $p < .001$) or outside home ($X^2 = 21.5$, $p < .001$), lower percentage of adolescents with screen use > 2 hs ($X^2 = 8.87$, $p = .003$), lower percentage of adolescents that assist to extra-curricular intellectual ($X^2 = 30.8$, $p < .001$) or physical activities ($X^2 = 12.8$, $p < .001$) (Table 1, statistics not shown in the table).

Table 2 shows the univariate and multivariate predictors of sleep deficit in adolescents without and with UBN. In those without UBN, sleep deficit was independently predicted by school distance, school starting time, full-day school, and working outside home. In those with UBN, sleep deficit was independently predicted by school starting time, full-day school, screen use > 2 hs and working at home, while being outdoor playing > 2 hs a protective factor. In both conditions, morning school starting time highly augments the risk of sleep deficit (Table 2).

Discussion

The main result of this study is the finding of a global high prevalence of short sleeping time, a slight increase of sleep time in adolescents with UBN, and different patterns of wake activities that predict sleep deficit, depending on the presence of UBN. The prevalence rate of reduced sleep of 38% identified in the population of this study is comparable to the one reported by our group in four cities from Argentina (49%) (Perez-Chada et al., 2007), and by others groups in Santa María (54%) (Pereira, Moreno, & Louzada, 2014) and São Paulo (39%) (Leite Bernardo, Felden Pereira, Mazzilli Louzada, & D'Almeida, 2009) from Brazil. These rates were obtained employing a cut-point of 8 hours to define sleep deficit, but on the other side, day-time sleeping (naps) was excluded from the calculation.

The presence of UBN increased risetime and sleep duration. This contradicts previous literature, where a lower socioeconomic background is usually associated with less sleep duration and more sleep disruption in adolescents (Knutson & Lauderdale, 2009; Leite Bernardo et al., 2009). Socioeconomic demographics like income, educational level, and employment status are usually associated with more delayed, shorter duration, and less consistent sleep patterns (Marco et al., 2011). However, none of these studies focused in situations of extreme poverty. Among the factors associated with the presence of UBN that may justify these findings, the assistance to nearby schools probably explain the increased risetime and indirectly the increased sleep time. The association between UBN and attendance

to neighborhood schools is as expected, since the seek for better schools tends to be available for families with higher socio-economic status through residential mobility and enrolment in private schools (Musset, 2012). Another factor that could explain the increased risetime is the observed lower percentage of children that assist to extra-curricular intellectual or physical activities. However, if this had been the case, those activities would have been mostly developed in the morning, and differences in the percentage of school morning assistance should have also been observed.

School starting time and full-day schooling were strong predictors of sleep deficit in adolescents with and without UBN. Starting school at the morning school is a well-recognized risk factor for sleep deprivation, determining less time spent in bed, worse sleep quality and increased daytime sleepiness which in turn leads to bad mood and poor performance (Wolfson & Carskadon, 1998; De Souza, Cortez de Sousa, Leao Maia, & Macedo de Azevedo, 2011; Perez-Lloret et al., 2013). Unlike other school systems, in Argentina some schools have half-day schedules while others full-day schedules. It is expected that the extended day pose a higher risk of sleep deficit, because it combines an early school starting with being at school most of the day (Kalenkoski, Ribar, & Stratton, 2009; Alasker & Flammer, 1998), thus preventing the possibility of taking naps or delaying bed time. As discussed below, this is more serious when adolescents are involved in some kind of work. In addition, in adolescents without UBN, school distance was a risk factor for sleep deficit, as reported previously by other authors. For example, in a sample of 1126 adolescents in a urban area from Brazil, longer journeys to school were associated with an earlier rise time and delay at the mid-sleep phase, especially in adolescents who started school at the morning, which led to a significantly reduced time in bed (Pereira et al., 2014).

Child labour appeared as a predictor of sleep deficit. In adolescents without UBN, the risk is associated with paid work, while in adolescents with UBN is associated with intensive work at home. Our results are consistent with those of previously published studies that show that adolescents who work wake up earlier and have decreased night and total sleep duration during the week than nonworking students (Teixeira et al., 2006; Vinha D et al., 2010; Fischer et al., 2008). Less information was found regarding the potentially sleep damaging effects of work at home. In a cross-national research comprising 2315 adolescents from Europe and United States, household work was negatively correlated with sleep and it was suggested that adolescents from poorer families were more involved in this kind of work (Alasker & Flammer, 1998). In this regard, maternal education correlates negatively with teens' time spent in

housework, while teenagers in households with higher incomes spend more time in paid work (Wight, Price, Bianchi, & Hunt, 2009).

Finally, it is worth describing the impact of other activities in sleep deficit. In adolescents with UBN, sleep deficit was independently predicted by screen use > 2hs, while being outdoor playing > 2hs a protective factor. Watching TV or using computers was found to delay bed time, decrease sleep duration and reduce subjective sleep quality (Knutson & Lauderdale, 2009; Mesquita & Reimao, 2007). Interestingly, TV use is increased in children from households with lower socioeconomic status (Stamatakis, Hillsdon, Mishra, Hamer, & Marmot, 2009) and computer use is increased in families with higher family incomes (Wight et al., 2009). Regarding the protective effect of playing outdoors, it could be understood as way of preventing excessive screen use and promoting sleep by several physiological mechanisms including sunlight exposure, thus representing a sleep hygiene measure (Buman & King, 2010).

Several limitations of this study must be taken into account. First, objective measures were not performed. Second, information about sleep schedules was provided by the parent or legal guardian, and not directly from the adolescent. Third, information about daily activities was gathered in the context of a social study not specifically aimed to assess factors that predict sleep. Nevertheless, we think that the information provided by this work is still valuable and highlights the utility of using nation-wide social surveys or census data to develop chronobiological studies.

To conclude, this study focused on the factors that may be involved in sleep deprivation of adolescents with UBN. Household work appeared as a risk factor that may increase the burden of sleep deficit, while playing outdoors appeared as a protective factor, adding to the well recognized risk factors of starting school early and watching screens for long periods. The reduced academic achievement, increased risk of accidents and adverse health outcomes associated with sleep deprivation poses sleep as an additional unsatisfied basic need that worsens living conditions of this group of teens. In this regard, our results may help to design public health policies that contribute to ameliorate this adverse situation.

Acknowledgments

Studies were supported by grants from the Agencia Nacional de Promoción Científica y Tecnológica, Argentina (PICT 2010-2195, PICT 2012-0984, PICT Start-Up 2013-0710). DEV, DPC and DG are Research Career Awardees from the Argentine Research Council (CONICET). Data used herein were provided by the Argentine Social Debt Barometer (Barómetro de la Deuda Social Argentina) through the Argentine Social Debt Survey (Encuesta sobre la Deuda Social Argentina).

References

Alasker, F. & Flammer, A. (1998). Time use by adolescents in an international perspective. II: The case of necessary activities. In A. Flammer & F. Alasker (Eds.), *The Adolescent Experience: European and American Adolescents in the 1990s* (pp. 61-84). Mahwah, NJ: Lawrence Erlbaum Associates.

Buman, M. & King, A. (2010). Exercise as a Treatment to Enhance Sleep. *American Journal of Lifestyle Medicine*, 4, 500-514.

Cappuccio, F. P., Taggart, F. M., Kandala, N. B., Currie, A., Peile, E., Stranges, S. et al. (2008). Meta-analysis of short sleep duration and obesity in children and adults. *Sleep*, 31, 619-626.

Cardinali, D., Simonelli, G., Rodríguez Espínola, S., Salvia, A., Pérez Chada, D., & Vigo, D. (2013). Sleep, Slums and Shelter: Impact of a Slum-housing Upgrading Program. In *Proceedings of the Working Group in Bread, Brain, Education and Poverty* (pp. 1-14). Vatican City: Pontifical Academy of Sciences. Available at: www.pas.va/content/dam/accademia/pdf/sv125/sv125-cardinali.pdf

Carskadon, M. A. (2011). Sleep in adolescents: the perfect storm. *Pediatr.Clin.North Am.*, 58, 637-647.

Carskadon, M. A., Acebo, C., & Jenni, O. G. (2004). Regulation of adolescent sleep: implications for behavior. *Ann.N.Y.Acad.Sci.*, 1021, 276-291.

De Souza, J., Cortez de Sousa, I., Leao Maia, A., & Macedo de Azevedo, C. (2011). Sleep patterns of teachers and adolescents who attend school in the morning. *Biological Rhythm Research*, 43, 65-72.

Fischer, F. M., Nagai, R., & Teixeira, L. R. (2008). Explaining sleep duration in adolescents: the impact of socio-demographic and lifestyle factors and working status. *Chronobiol.Int.*, 25, 359-372.

International Labour Office - International Programme on the Elimination of Child Labour (IPEC) (2013). *Marking progress against child labour: Global estimates and trends 2000-2012*. Geneva: ILO.

Kalenkoski, C., Ribar, D., & Stratton, L. (2009). How do adolescents spend time use? *IZA Discussion Papers*, 4374.

Knutson, K. L. & Lauderdale, D. S. (2009). Sociodemographic and behavioral predictors of bed time and wake time among US adolescents aged 15 to 17 years. *J.Pediatr.*, 154, 426-30, 430.

Leite Bernardo, M., Felden Pereira, E., Mazzilli Louzada, F., & D'Almeida, V. (2009). Duração do sono em adolescentes de diferentes níveis socioeconômicos. *J Bras Psiquiatr*, 58, 231-237.

Marco, C. A., Wolfson, A. R., Sparling, M., & Azuaje, A. (2011). Family socioeconomic status and sleep patterns of young adolescents. *Behav.Sleep Med.*, 10, 70-80.

Mesquita, G. & Reimao, R. (2007). Nightly use of computer by adolescents: its effect on quality of sleep. *Arq Neuropsiquiatr.*, 65, 428-432.

Mindell, J. A. & Meltzer, L. J. (2008). Behavioural sleep disorders in children and adolescents. *Ann.Acad.Med.Singapore*, 37, 722-728.

Musset, P. (2012). School Choice and Equity: Current Policies in OECD Countries and a Literature Review. *OECD Education Working Papers*, 66.

Patel, N. P., Grandner, M. A., Xie, D., Branas, C. C., & Gooneratne, N. (2010). "Sleep disparity" in the population: poor sleep quality is strongly associated with poverty and ethnicity. *BMC.Public Health*, 10, 475.

Pereira, E. F., Moreno, C., & Louzada, F. M. (2014). Increased commuting to school time reduces sleep duration in adolescents. *Chronobiol.Int.*, 31, 87-94.

Perez-Chada, D., Drake, C. L., Perez-Lloret, S., Videla, A. J., & Cardinali, D. P. (2009). Brief review: diurnal rhythms, obesity and educational achievement in South american cultures. *Int.J.Neurosci.*, 119, 1091-1104.

Perez-Chada, D., Perez-Lloret, S., Videla, A. J., Cardinali, D., Bergna, M. A., Fernandez-Acquier, M. et al. (2007). Sleep disordered breathing and daytime sleepiness are associated with poor academic performance in teenagers. A study using the Pediatric Daytime Sleepiness Scale (PDSS). *Sleep*, 30, 1698-1703.

Perez-Lloret, S., Videla, A. J., Richaudeau, A., Vigo, D., Rossi, M., Cardinali, D. P. et al. (2013). A multi-step pathway connecting short sleep duration to daytime somnolence, reduced attention, and poor academic performance: an exploratory cross-sectional study in teenagers. *J.Clin.Sleep Med.*, 9, 469-473.

Roberts, R., Roberts, C., & Chen, I. (2001). Functioning of adolescents with symptoms of disturbed sleep. *Journal of Youth and Adolescence*, 30, 1-18.

Simonelli, G., Leanza, Y., Boilard, A., Hyland, M., Augustinavicius, J. L., Cardinali, D. P. et al. (2013). Sleep and quality of life in urban poverty: the effect of a slum housing upgrading program. *Sleep*, 36, 1669-1676.

Stamatakis, E., Hillsdon, M., Mishra, G., Hamer, M., & Marmot, M. (2009). Television viewing and other screen-based entertainment in relation to multiple socioeconomic status indicators and area deprivation: the Scottish Health Survey 2003. *J.Epidemiol.Community Health*, 63, 734-740.

Teixeira, L. R., Fischer, F. M., & Lowden, A. (2006). Sleep deprivation of working adolescents--a hidden work hazard. *Scand.J.Work Environ.Health*, 32, 328-330.

Tuñón, I. (2013). *Hacia el pleno ejercicio de derechos en la niñez y adolescencia: propensiones, retos y desigualdades en la Argentina urbana (2010-2012)*. Buenos Aires: Educa.

Vinha D, Cavalcante JA, & Andrade M (2010). Sleep-Wake Patterns of Student Workers and Non-Workers. *Biological Rhythm Research*, 33, 417-426.

Wight, V., Price, J., Bianchi, S., & Hunt, B. (2009). The time use of teenagers. *Social Science Research*, 38, 792-809.

Wolfson, A. R. & Carskadon, M. A. (1998). Sleep schedules and daytime functioning in adolescents. *Child Dev.*, 69, 875-887.

Wolfson, A. R. & Carskadon, M. A. (2003). Understanding adolescents' sleep patterns and school performance: a critical appraisal. *Sleep Med.Rev*, 7, 491-506.

Table 1: Unsatisfied basic needs and sleep deficit

	Unsatisfied Basic Needs								
	No (n=1292, 77%)			Yes (n=390, 23%)			Total (n=1682, 100%)		
	Sleep deficit			Sleep deficit			Sleep deficit		
	No (n=700, 61%)	Yes (n=454, 39%)	Total (n=1154, 100%)	No (n=257, 67%)	Yes (n=125, 33%)	Total (n=382, 100%)	No (n=957, 62%)	Yes (n=579, 38%)	Total (n=1536, 100%)
Demographics									
Female	333 (48%)	225 (50%)	558 (48%)	124 (48%)	65 (52%)	189 (49%)	457 (48%)	290 (50%)	747 (49%)
Age (ys)	14.9 ± .05	15.0 ± .06	15.0 ± .04	15.0 ± .08	14.9 ± .11	15.0 ± .07	15.0 ± .04	15.0 ± .06	15.0 ± .03
Sleep									
Bed time (hs)	22.8 ± .05	23.4 ± .04	23.1 ± .03	22.7 ± .07	23.4 ± .07	22.9 ± .06	22.8 ± .04	23.4 ± .04	23.0 ± .03
Rise time (hs)	8.6 ± .06	6.8 ± .03	7.9 ± .05	8.9 ± .10	6.8 ± .04	8.2 ± .08	8.7 ± .05	6.8 ± .03	8.0 ± .04
Night sleep (hs)	9.8 ± .05	7.4 ± .04	8.8 ± .05	10.2 ± .09	7.4 ± .06	9.3 ± .09	9.9 ± .04	7.4 ± .03	8.9 ± .04
Nap habit	164 (23%)	66 (14%)	229 (20%)	54 (21%)	13 (10%)	66 (17%)	218 (23%)	78 (14%)	296 (19%)
Nap duration (hs)	1.8 ± .06	1.4 ± .07	1.7 ± .05	2.0 ± .12	1.3 ± .13	1.9 ± .10	1.9 ± .06	1.3 ± .06	1.7 ± .05
TST (hs)	10.2 ± .05	7.6 ± .04	9.2 ± .05	10.6 ± .09	7.6 ± .06	9.6 ± .10	10.3 ± .04	7.6 ± .03	9.3 ± .04
School									
Distance > 1km	229 (37%)	220 (52%)	449 (43%)	57 (27%)	36 (30%)	93 (28%)	285 (34%)	256 (47%)	541 (39%)
Morning	302 (48%)	315 (71%)	617 (58%)	91 (43%)	96 (81%)	187 (57%)	393 (47%)	411 (73%)	804 (57%)
Afternoon	280 (45%)	40 (9%)	320 (30%)	106 (50%)	9 (8%)	116 (35%)	386 (46%)	49 (9%)	436 (31%)
Evening	9 (1%)	6 (1%)	15 (1%)	5 (2%)	0 (0%)	5 (2%)	14 (2%)	6 (1%)	20 (1%)
Full-day	37 (6%)	80 (18%)	117 (11%)	10 (5%)	14 (11%)	24 (7%)	47 (6%)	93 (17%)	140 (10%)
Child labour									
Home work	70 (10%)	33 (7%)	103 (9%)	62 (24%)	35 (28%)	97 (25%)	132 (14%)	69 (12%)	201 (13%)
Paid work	104 (15%)	76 (17%)	180 (16%)	64 (25%)	33 (26%)	97 (25%)	168 (18%)	109 (19%)	277 (18%)
Other activities									
Screen >2hs	473 (68%)	316 (70%)	789 (68%)	141 (55%)	88 (70%)	229 (60%)	614 (64%)	404 (70%)	1019 (66%)
Outdoor > 2hs	206 (33%)	133 (32%)	339 (33%)	86 (40%)	49 (42%)	135 (40%)	291(35%)	183 (34%)	474 (35%)
Extra-curricular intellectual	186 (27%)	173 (38%)	359 (31%)	41 (16%)	29 (23%)	70 (18%)	227 (24%)	202 (35%)	429 (28%)
Extra-curricular physical	308 (44%)	250 (55%)	558 (48%)	101 (39%)	46 (37%)	147 (38%)	409 (43%)	296 (51%)	705 (46%)

Values are expressed as mean ± standard error for numerical variables or frequency (percentage) for categorical variables. TST: total sleep time.

Table 2: Univariate and multivariate predictors of sleep deficit in adolescents

	UBN = no				UBN = yes			
	Univariate (1)		Multivariate (2)		Univariate (1)		Multivariate (2)	
	T or X ²	p	OR (95% CI)	p	T or X ²	p	OR (95% CI)	p
Demographics								
Female	.41	.524	-	-	.52	.470	-	-
Age (ys)	-1.28	.202	-	-	1.35	.179	-	-
School								
Distance > 1km	23.9	<.001	1.77 (1.33-2.36)	<.001	.53	.467	-	-
Morning	133	<.001	8.35 (5.66-12.31)	<.001	59	<.001	14.45 (6.76-30.89)	<.001
Full-day	134	<.001	14.0 (8.1-24.5)	<.001	37	<.001	18.39 (5.82-58.11)	<.001
Child labour								
Home work	2.5	.112	-	-	.67	.414	2.14 (1.14-4.03)	.019
Paid work	.77	.380	1.67 (1.10-2.55)	.017	.113	.737	-	-
Other activities								
Screen >2hs	.53	.47	-	-	8.45	.004	2.15 (1.23-3.76)	.007
Outdoor >2hs	.06	.806	-	-	.11	.736	.38 (.18-.80)	.011
Extra-curricular intellectual	17.0	<.001	-	-	2.95	.086	-	-
Extra-curricular physical	13.5	<.001	-	-	.28	.598	-	-

OR: Odds ratio. 95% CI: 95% confidence interval. TST: total sleep time; UBN: Unsatisfied basic needs. (1) Univariate analyses: T test (numerical variables) or chi-square test (categorical variables). (2) Multivariate analyses: Binary logistic regression. Model summary without UBN: 82.2% cases included, 68.9% cases predicted, $R^2 = 0.248$; with UBN: 80.6% cases included, 72.6% predicted. $R^2 = 0.345$.