

**Fustinori, Osvaldo ; Saposnik, Gustavo ; Esnaola y Rojas, María Martha ; Lakkis, S. G. ; Sposato, Luciano A.**

*Higher frequency of atrial fibrillation linked to colder seasons and air temperature on the day of ischemic stroke onset*

**Documento publicado en Journal of Stroke and Cerebrovascular Diseases, 22 (4) 2013**

Este documento está disponible en la Biblioteca Digital de la Universidad Católica Argentina, repositorio institucional desarrollado por la Biblioteca Central “San Benito Abad”. Su objetivo es difundir y preservar la producción intelectual de la Institución.

La Biblioteca posee la autorización del autor para su divulgación en línea.

Cómo citar el documento:

Fustinori, O., G. Saposnik, M. M. Esnaola y Rojas, S. G. Lakkis y L. A. Sposato. 2013. Higher frequency of atrial fibrillation linked to colder seasons and air temperature on the day of ischemic stroke onset [en línea]. *Journal of stroke and cerebrovascular diseases* 22(4). <http://www.sciencedirect.com/science/article/pii/S1052305713000748>  
Disponible en: <http://bibliotecadigital.uca.edu.ar/repositorio/investigacion/higher-frequency-atrial-fibrillation.pdf>  
[Fecha de consulta: ....]

# Higher Frequency of Atrial Fibrillation Linked to Colder Seasons and Air Temperature on the Day of Ischemic Stroke Onset

Osvaldo Fustinoni, MD,\* Gustavo Saposnik, MD, MSc, FAHA,<sup>†‡§||¶</sup>  
 María Martha Esnaola y Rojas, MD,# Susan G. Lakkis, PhD,\*\* and  
 Luciano A. Sposato, MD, MBA, FAHA,<sup>†††</sup> On behalf of ReNACer Investigators

**Background:** Whether a seasonal variation of atrial fibrillation among acute ischemic stroke (AIS) patients occurs is unknown. We studied the distribution of atrial fibrillation across seasons and air temperatures in a cohort of AIS patients. **Methods:** We selected 899 AIS patients from the Argentinean Stroke Registry (ReNACer), who were admitted to 43 centers in the Province of Buenos Aires. We recorded the minimum and maximum temperatures at local weather centers on the day and the city where each stroke occurred. We used the goodness-of-fit  $\chi^2$  test to assess the distribution of atrial fibrillation across seasons and air temperatures and the Pearson correlation coefficient to assess the relationship between these variables. We developed a regression model for testing the association between seasons and atrial fibrillation. **Results:** We found a seasonal variation in the occurrence of atrial fibrillation, with a peak in winter and a valley in summer (23.1% versus 14.0%,  $P < .001$ ). The semester comprised by autumn and winter was associated with atrial fibrillation (Pearson  $P < .001$ ). Atrial fibrillation showed a nonhomogeneous distribution across ranges of temperature ( $P < .001$ , goodness-of-fit test), with a peak between 5°C and 9°C, and was associated with minimum (Pearson  $P = .042$ ) and maximum (Pearson  $P = .002$ ) air temperature. After adjusting for significant covariates, there was a 2-fold risk of atrial fibrillation during autumn and winter. **Conclusions:** In this cohort of AIS patients, atrial fibrillation showed a seasonal variation and a nonhomogeneous distribution across air temperatures, with peaks in cold seasons and low temperatures on the day of stroke onset. **Key Words:** Ischemic stroke—atrial fibrillation—risk—season—temperature—weather.

© 2013 by National Stroke Association

From the \*Cerebrovascular Diseases, Instituto de Neurociencias Buenos Aires, Buenos Aires, Argentina; †Stroke Outcomes Research Center, St Michael's Hospital, Ontario, Canada; ‡Li Ka Shing Knowledge Institute, St Michael's Hospital, Ontario, Canada; §Department of Medicine, University of Toronto, Ontario, Canada; ||Department of Health Policy Management and Evaluation, University of Toronto, Ontario, Canada; ¶Institute for Clinical Evaluative Sciences, Ontario, Canada; #Department of Neurology, Hospital Dr Cesar Milstein, Buenos Aires, Argentina; \*\*Equipo Interdisciplinario para el Estudio del Cambio Global, Pontificia Universidad Católica Argentina, Puerto Madero, Argentina; ††Stroke Center at the Institute of Neurosciences, University Hospital, Favaloro Foundation, Buenos Aires, Argentina;

and ‡‡Vascular Research Unit, INECO Foundation, Buenos Aires, Argentina.

Received February 7, 2013; revision received March 6, 2013; accepted March 7, 2013.

Address correspondence to Luciano A. Sposato, MD, MBA, FAHA, Stroke Center at the Institute of Neurosciences, University Hospital, Favaloro Foundation and Vascular Research Unit, INECO Foundation, Rosales 2616, Olivos, Buenos Aires, Argentina. E-mail: lucianospasato@gmail.com.

1052-3057/\$ - see front matter

© 2013 by National Stroke Association

<http://dx.doi.org/10.1016/j.jstrokecerebrovasdis.2013.03.009>